Interdistrict School Choice: A Theory of Student Assignment

Authors: Isa Hafalir, Fohita Kojima, M. Bumin Yenmez

Persistent link: http://hdl.handle.net/2345/bc-ir:108266

This work is posted on eScholarship@BC, Boston College University Libraries.

Boston College Working Papers in Economics, 2018

Originally posted on: http://ideas.repec.org/p/boc/bocoec/970.html

INTERDISTRICT SCHOOL CHOICE: A THEORY OF STUDENT ASSIGNMENT[†]

ISA E. HAFALIR, FUHITO KOJIMA, AND M. BUMIN YENMEZ*

ABSTRACT. Interdistrict school choice programs—where a student can be assigned to a school outside of her district—are widespread in the US, yet the market-design literature has not considered such programs. We introduce a model of interdistrict school choice and present two mechanisms that produce *stable* or *efficient* assignments. We consider three categories of policy goals on assignments and identify when the mechanisms can achieve them. By introducing a novel framework of interdistrict school choice, we provide a new avenue of research in market design.

1. Introduction

School choice is a program that uses preferences of children and their parents over public schools to assign children to schools. It has expanded rapidly in the United States and many other countries in the last few decades. Growing popularity and interest in school choice stimulated research in market design, which has not only studied this problem in the abstract, but also contributed to designing specific assignment mechanisms.¹

Existing market-design research about school choice is, however, limited to *intradistrict* choice, where each student is assigned to a school only in her own district. In other words, the literature has not studied *interdistrict* choice, where a student can be assigned to a school outside of her district. This is a severe limitation for at least two reasons. First, interdistrict school choice is widespread: some form of it is practiced in 43 U.S. states.² Second, as we illustrate in detail below, many policy goals in school choice impose constraints across districts in reality, but the existing literature assumes away such constraints. This omission limits our ability to analyze these policies of interest.

Date: December 28, 2018, First draft: July 15, 2017.

Keywords: Interdistrict school choice, student assignment, stability, efficiency.

We thank Haluk Ergin, Yuichiro Kamada, Tayfun Sönmez, Utku Ünver, Rakesh Vohra, and the audiences at various seminars and conferences. Lukas Bolte, Ye Rin Kang, and especially Kevin Li provided superb research assistance. Kojima acknowledges financial support from the National Research Foundation through its Global Research Network Grant (NRF-2016S1A2A2912564). Hafalir is affiliated with the UTS Business School, University of Technology Sydney, Sydney, Australia; Kojima is with the Department of Economics, Stanford University, 579 Serra Mall, Stanford, CA, 94305; Yenmez is with the Department of Economics, Boston College, 140 Commonwealth Ave, Chestnut Hill, MA, 02467. Emails: isa.hafalir@uts.edu.au, fkojima@stanford.edu, bumin.yenmez@bc.edu.

¹See Abdulkadiroğlu et al. (2005a,b, 2009) for details of the implementation of these new school choice procedures in New York and Boston.

²See http://ecs.force.com/mbdata/mbquest4e?rep=0E1705, accessed on July 14, 2017.

In this paper, we propose a model of interdistrict school choice. Our paper builds upon matching models in the tradition of Gale and Shapley (1962).³ We study algorithms and interdistrict admissions rules to assign students to schools under which a variety of policy goals can be established, an approach similar to the intradistrict school choice literature (Abdulkadiroğlu and Sönmez, 2003). In our setting, however, policy goals are defined on the district level—or sometimes even over multiple districts—rather than the individual school level, placing our model outside of the standard setting. To facilitate the analysis in this setting, we model the problem as matching with contracts (Hatfield and Milgrom, 2005) between students and districts in which a contract specifies the particular school within the district that the student attends.⁴

Following the school choice literature, we begin our analysis by considering *stability* (we also consider efficiency, as explained later). To define stability in our framework, we assume that each district is endowed with an admissions rule represented by a choice function over sets of contracts. We focus our attention on the student-proposing deferred-acceptance mechanism (SPDA) of Gale and Shapley (1962). In our setting, this mechanism is not only stable but also strategy-proof—i.e., it renders truthtelling a weakly dominant strategy for each student.

In this context, we formalize a number of important policy goals. The first is *individual rationality* in the sense that every student is matched with a weakly more preferred school than the school she is initially matched with (in the absence of interdistrict school choice). This is an important requirement, because if an interdistrict school choice program harms students, then public opposition is expected and the program may not be sustainable. The second policy is what we call *the balanced-exchange policy*, that the number of students that each district receives from the other districts must be the same as the number of students that it sends to the others. Balanced exchange is also highly desired by school districts in practice. This is because each district's funding depends on the number of students that it serves and, therefore, if the balanced-exchange policy is not satisfied, then some districts may lose funding, possibly making the interdistrict school choice program impossible. For each of these policy goals, we identify the necessary and sufficient condition for achieving that goal under SPDA as a restriction on district admissions rules.

Last, but not least, we also consider a requirement that there be enough student diversity in each district. In fact, diversity appears to be the main motivation for many interdistrict school choice programs.⁵ To put this into context, we note that the lack of diversity is

³We use the terms *assignment* and *matching* interchangeably for the rest of the paper.

⁴One might suspect that an interdistrict school choice problem can readily be reduced to an intradistrict problem by relabeling a district as a school. This is not the case because, among other things, which school within a district a student is matched with matters for that student's welfare.

⁵We refer to Wells et al. (2009) for a review and discussion of interdistrict integration programs.

prevalent under intradistrict school choice programs even though they often seek diversity by controlled-choice constraints.⁶ This is perhaps unsurprising given that only residents of the given district can participate in intradistrict school choice and there is often severe residential segregation. In fact, a number of studies such as Rivkin (1994) and Clotfelter (1999, 2011) attribute the majority—as high as 80 percent for some data and measure of racial and ethnic segregation in public schools to disparities between school districts rather than within school districts. Given this concern, many interdistrict choice programs explicitly list achieving diversity as their main goal.

A case in point is the *Achievement and Integration (AI) Program* of the Minnesota Department of Education (MDE). Introduced in 2013, the AI program incentivizes school districts for integration. A district is required to participate in this program if the proportion of a minority group in the district is considerably higher than that in a neighboring district. In particular, every year the MDE commissioner analyzes fall enrollment data from every district and, when a district and one of its adjoining districts have a difference of 20 percent or higher in the proportion of any group of enrolled *protected students* (American Indian, Asian or Pacific Islander, Hispanic, Black, not of Hispanic origin, and White, not of Hispanic origin), the district with the higher percentage is required to be in the AI program.⁷ In the 2015-16 school year, more than 120 school districts participated in this program (Figure 1, taken from MDE's website, shows school districts in the Minneapolis-Saint Paul metro area that take part in this program).

Motivated by Minnesota's AI program, we consider a policy goal requiring that the difference in the proportions of each student type across districts be within a given bound. Then, we provide a necessary and sufficient condition for SPDA to satisfy the diversity policy. The condition provided is one on district admissions rules that have a structure of type-specific ceilings, an analogue of the class of choice rules analyzed by Abdulkadiroğlu and Sönmez (2003) and Ehlers et al. (2014) in the context of a more standard intradistrict school-choice problem.

Next, we turn our attention to efficiency. Given that the distributional policy goals work as constraints on matchings, we use the concept of *constrained efficiency*. We say that a matching is constrained efficient if it satisfies the policy goal and is not Pareto dominated by any matching that satisfies the same policy goal. In addition, we require individual

⁶Examples of controlled school choice include Boston before 1999, Cambridge, Columbus, and Minneapolis. See Abdulkadiroğlu and Sönmez (2003) for details of these programs as well as analysis of controlled school choice.

⁷In Minnesota's AI program, if the difference in the proportion of protected students at a school is 20 percent or higher than a school in the same district, the school with the higher percentage is considered a *racially identifiable school* (RIS) and districts with RIS schools also need to participate in the AI program. In this paper, we focus on diversity issues across districts rather than within districts. Diversity problems within districts are studied in the controlled school choice literature that we discuss below.



FIGURE 1. Minnesota-Saint Paul metro area school districts participating in the AI program. The districts with the same color are adjoining districts that exchange students with one another.

rationality and strategy-proofness.⁸ We first demonstrate an impossibility result; when the diversity policy is given as type-specific ceilings at the district level, there is no mechanism that satisfies the policy goal, constrained efficiency, individual rationality, and strategy-proofness. By contrast, a version of the top trading cycles (TTC) mechanism (Shapley and Scarf, 1974) satisfies these properties when the policy goal satisfies M-convexity, a concept in discrete mathematics (Murota, 2003). We proceed to show that the balanced-exchange policy and an alternative form of diversity policy—type-specific ceilings at the individual school level instead of at the district level—are M-convex, so TTC satisfies the desired properties for these policies. The same conclusion holds even when both of these policy goals are imposed simultaneously.

⁸Without individual rationality, all the other desired properties can be attained by a serial dictatorship.

We also consider the case when there is a policy function that measures how well a matching satisfies the policy goal. For example, diversity of a matching can be measured as its distance to an ideal distribution of students. We show that TTC satisfies the same desirable properties when the policy function satisfies *pseudo M-concavity*, a notion of concavity for discrete functions that we introduce. Furthermore, we show that there is an equivalence between two approaches based on the M-convexity of the policy set and the pseudo M-concavity of the policy function. Therefore, both results can naturally be applied in different settings depending on how the policy goals are stated.

Related Literature. Our paper is closely related to the controlled school choice literature that studies student diversity in schools in a given district. Abdulkadiroğlu and Sönmez (2003) introduce a policy that imposes type-specific ceilings on each school. This policy has been analyzed by Abdulkadiroğlu (2005), Ergin and Sönmez (2006), and Kojima (2012), among others. More accommodating policies using reserves rather than type-specific ceilings have been proposed and analyzed by Hafalir et al. (2013) and Ehlers et al. (2014). The latter paper finds difficulties associated with hard floor constraints, an issue further analyzed by Fragiadakis et al. (2015) and Fragiadakis and Troyan (2017).⁹ In addition to sharing the motivation of achieving diversity, our paper is related to this literature in that we extend the type-specific reserve and ceiling constraints to district admissions rules. In contrast to this literature, however, our policy goals are imposed on districts rather than individual schools, which makes our model and analysis different from the existing ones.

The feature of our paper that imposes constraints on sets of schools (i.e., districts), rather than individual schools, is shared by several recent studies in matching with constraints. Kamada and Kojima (2015) study a model where the number of doctors who can be matched with hospitals in each region has an upper bound constraint. Variations and generalizations of this problem are studied by Goto et al. (2014, 2017), Biro et al. (2010), and Kamada and Kojima (2017, 2018), among others. While sharing the broad interest in constraints, these papers are different from ours in at least two major respects. First, they do not assume a set of hospitals is endowed with a well-defined choice function, while each school district has a choice function in our model. Second, the policy issues studied in these papers and those studied in ours are different given differences in the intended applications. These differences render our analysis distinct from those of the other papers, with none of their results implying ours and vice versa.

One of the notable features of our model is that district admissions rules do not necessarily satisfy the standard assumptions in the literature, such as *substitutability*, which

⁹In addition to the works discussed above, recent studies on controlled school choice and other twosided matching problems with diversity concerns include Westkamp (2013), Echenique and Yenmez (2015), Sönmez (2013), Kominers and Sönmez (2016), Dur et al. (2014), Dur et al. (2016), and Nguyen and Vohra (2017).

guarantee the existence of a stable matching. In fact, even a seemingly reasonable district admissions rule may violate substitutability because a district can choose at most one contract associated with the same student—namely just one contract representing one school that the student can attend. Rather, we make weaker assumptions following the approach of Hatfield and Kominers (2014). This issue is playing an increasingly prominent role in matching with contracts literature; for example, in matching with constraints (Kamada and Kojima, 2015), college admissions (Aygün and Turhan, 2016; Yenmez, 2018), and post-graduate admissions (Hassidim et al., 2017), to name just a few.

Our analysis of Pareto efficient mechanisms is related to a small but rapidly growing literature that uses discrete optimization techniques for matching problems. Closest to ours is Suzuki et al. (2017), who show that a version of the TTC mechanism satisfies desirable properties if the constraint satisfies M-convexity.¹⁰ Our analysis on efficiency builds upon and generalizes theirs. While the use of discrete convexity concepts for studying efficient object allocation is still rare, it has been utilized in an increasing number of matching problems such as two-sided matching with possibly bounded transfer (Fujishige and Tamura, 2006, 2007), matching with substitutable choice functions (Murota and Yokoi, 2015), matching with constraints (Kojima et al., 2018a), and trading networks (Candogan et al., 2016).

There is also a recent literature on segmented matching markets in a given district. Manjunath and Turhan (2016) study a setting where different clearinghouses can be coordinated, but not integrated, in a centralized clearinghouse and show how a stable matching can be achieved. In a similar setting, Dur and Kesten (2018) study sequential mechanisms and show that these mechanisms lack desired properties. In another work, Ekmekci and Yenmez (2014) study the incentives of a school to join a centralized clearinghouse. In contrast to these papers, we study which interdistrict school choice policies can be achieved when districts are integrated.

At a high level, the present paper is part of research in resource allocation under constraints. Real-life auction problems often feature constraints (Milgrom, 2009), and a great deal of attention was paid to cope with complex constraints in a recent FCC auction for spectrum allocation (Milgrom and Segal, 2014). Auction and exchange markets under constraints are analyzed by Bing et al. (2004), Gul et al. (2018), and Kojima et al. (2018b). Handling constraints is also a subject of a series of papers on probabilistic assignment mechanisms (Budish et al., 2013; Che et al., 2013; Pycia and Ünver, 2015; Akbarpour and Nikzad, 2017; Nguyen et al., 2016). Closer to ours are Dur and Ünver (2018) and Dur et al. (2015). They consider the balance of incoming and outgoing members—a requirement

¹⁰See Kurata et al. (2016) for an earlier work on TTC in a more specialized setting involving floor constraints at individual schools.

that we also analyze—while modeling exchanges of members of different institutions under constraints. Although the differences in the model primitives and exact constraints make it impossible to directly compare their studies with ours, these papers and ours clearly share broad interests in designing mechanisms under constraints.

The rest of the paper is organized as follows. Section 2 introduces the model. In Sections 3 and 4, we study when the policy goals can be satisfied together with stability and constrained efficiency, respectively. Section 5 concludes. Additional results, examples, and omitted proofs are presented in the Appendix.

2. Model

In this section, we introduce our concepts and notation.

2.1. Preliminary Definitions. There exist finite sets of students S, districts D, and schools C. Each student s and school c has a home district denoted by d(s) and d(c), respectively. Each student s has a type $\tau(s)$ that can represent different aspects of the student such as the gender, race, socioeconomic status, etc. The set of all types is finite and denoted by T. Each school c has a capacity q_c , which is the maximum number of students that the school can enroll. There exist at least two school districts with one or more schools. For each district d, k_d is the number of students whose home district is d. In each district, schools have sufficiently large capacities to accommodate all students from the district, i.e., for every district d, $k_d \leq \sum_{c:d(c)=d} q_c$. For each type t, k^t is the number of type-t students.

Throughout this paper, we model interdistrict school choice as a matching problem between students and districts. However, merely identifying the district with which a student is matched leaves the specific school she is enrolled in unspecified. To specify which school within a district the student is matched with, we use the notion of contracts: A contract x = (s, d, c) specifies a student s, a district d, and a school c within this district, i.e., d(c) = d.¹¹ For any contract x, let s(x), d(x), and c(x) denote the student, district, and school associated with this contract, respectively. Let $\mathcal{X} \equiv \{(s, d, c) | d(c) = d\}$ denote the set of all contracts. For any set of contracts X, let X_s denote the set of contracts in X associated with student s, i.e., $X_s = \{x \in X | s(x) = s\}$. Similarly, let X_d and X_c denote the sets of contracts in X associated with district d and school c, respectively.

Each district *d* has an *admissions rule* that is represented by a choice function Ch_d . Given a set of contracts *X*, the district chooses a subset of contracts associated with itself, i.e., $Ch_d(X) = Ch_d(X_d) \subseteq X_d$.

Each student *s* has a strict preference order P_s over all schools and the outside option of being unmatched, which is denoted by \emptyset . Likewise, P_s is also used to rank contracts

¹¹For ease of exposition, a contract will sometimes be denoted by a pair (s, c) with the understanding that the district associated with the contract is the home district of school c.

associated with *s*. Furthermore, we assume that the outside option is the least preferred outcome, so for every contract *x* associated with *s*, *x* $P_s \emptyset$. The corresponding weak order is denoted by R_s . More precisely, for any two contracts *x*, *y* associated with *s*, *x* $R_s y$ if $x P_s y$ or x = y.

A *matching* is a set of contracts. A matching X is *feasible for students* if there exists at most one contract associated with every student in X. A matching X is *feasible* if it is feasible for students and the number of contracts associated with every school in X is at most its capacity, i.e., for any $c \in C$, $|X_c| \leq q_c$. We assume that there exists a feasible *initial matching* \tilde{X} such that every student has exactly one contract.¹² For any student s, if $\tilde{X}_s = \{(s, d, c)\}$ for some district d and school c, then c is called the *initial school* of s.

A *problem* is a tuple $(S, D, C, T, \{d(s), \tau(s), P_s\}_{s \in S}, \{Ch_d\}_{d \in D}, \{d(c), q_c\}_{c \in C}, \tilde{X}\}$. In what follows, we assume that all the components of a problem are publicly known except for student preferences. Therefore, we sometimes refer to a problem by the student preference profile which we denote as P_S . The preference profile of a subset of students $S \subseteq S$ is denoted by P_S .

2.2. Properties of Admissions Rules. A district admissions rule Ch_d is *feasible* if it always chooses a feasible matching. It is *acceptant* if, for any contract x associated with district d and matching X that is feasible for students; and if x is rejected from X, then at $Ch_d(X)$, either

- the number of students assigned to school c(x) is equal to $q_{c(x)}$, or
- the number of students assigned to district d is at least k_d .

In words, when a district admissions rule is acceptant, a contract x = (s, d, c) can be rejected by district d from a set which is feasible for students only if either the capacity of school c is filled or district d has accepted at least k_d students. Equivalently, if neither of these two conditions is satisfied, then the district has to accept the student. Throughout the paper, we assume that admissions rules are feasible and acceptant.¹³

A district admissions rule satisfies *substitutability* if, whenever a contract is chosen from a set, it is also chosen from any subset containing that contract (Kelso and Crawford, 1982; Roth, 1984). More formally, a district admissions rule Ch_d satisfies substitutability if, for every $x \in X \subseteq Y \subseteq \mathcal{X}$ with $x \in Ch_d(Y)$, it must be that $x \in Ch_d(X)$. A district admissions rule satisfies *the law of aggregate demand* (LAD) if the number of contracts chosen from a set is weakly greater than that of any of its subsets (Hatfield and Milgrom, 2005). Mathematically, a district admissions rule Ch_d satisfies LAD if, for every $X \subseteq Y \subseteq \mathcal{X}$,

¹²In Appendix A.1, we also consider the case when the initial matching for each district is constructed using student preferences and district admissions rules.

¹³In Section 3.3, we assume a weaker notion of acceptance when the admissions rule limits the number of students of each type that the district can accept.

 $|Ch_d(X)| \leq |Ch_d(Y)|$.¹⁴ A *completion* of a district admissions rule Ch_d is another admissions rule Ch'_d such that for every matching X either $Ch'_d(X)$ is equal to $Ch_d(X)$ or it is not feasible for students (Hatfield and Kominers, 2014). Throughout the paper, we assume that district admissions rules have completions that satisfy substitutability and LAD.¹⁵ In Appendix B, we provide classes of district admissions rules that satisfy our assumptions.

2.3. Matching Properties, Policy Goals, and Mechanisms. A feasible matching *X* satisfies *individual rationality* if every student weakly prefers her outcome in *X* to her initial school, i.e., for every student *s*, $X_s R_s \tilde{X}_s$.

A *distribution* $\xi \in \mathbb{Z}_{+}^{|\mathcal{C}| \times |\mathcal{T}|}$ is a vector such that the entry for school c and type t is denoted by ξ_c^t . The entry ξ_c^t is interpreted as the number of type-t students in school c at ξ . Furthermore, $\xi_d^t \equiv \sum_{c:d(c)=d} \xi_c^t$ denotes the number of type-t students in district d at ξ . Likewise, for any feasible matching X, the *distribution associated with* X is $\xi(X)$ whose c, t entry $\xi_c^t(X)$ is the number of type-t students assigned to school c at X. Similarly, $\xi_d^t(X)$ denotes the number of type-t students assigned to district d at X.

We represent a distributional policy goal as a set of distributions. Let Ξ denote a generic set of distributions. The policy that each student is matched without assigning any school more students than its capacity is denoted by Ξ^0 , i.e., $\Xi^0 \equiv \{\xi | \sum_{c,t} \xi_c^t = \sum_d k_d \text{ and } q_c \ge \sum_t \xi_c^t \text{ for all } c\}$. A matching *X* satisfies the policy goal Ξ if the distribution associated with *X* is in Ξ .

A feasible matching *X Pareto dominates* another feasible matching *Y* if every student weakly prefers her outcome in *X* to her outcome in *Y* and at least one student strictly prefers the former to the latter. Given a distributional policy goal, a feasible matching *X* that satisfies the policy goal satisfies *constrained efficiency* if there exists no feasible matching that satisfies the policy goal and Pareto dominates *X*.

A matching *X* is *stable* if it is feasible and

- districts would choose all contracts assigned to them, i.e., $Ch_d(X) = X_d$ for every district *d*, and
- there exist no student *s* and no district *d* who would like to match with each other, i.e., there exists no contract $x = (s, d, c) \notin X$ such that $x P_s X_s$ and $x \in Ch_d(X \cup \{x\})$.

Stability was introduced by Gale and Shapley (1962) for the college admissions problem. In the context of assigning students to public schools, it is viewed as a fairness notion (Abdulkadiroğlu and Sönmez, 2003).

A *mechanism* ϕ takes a profile of student preferences as input and produces a feasible matching. The outcome for student *s* at the reported preference profile P_S under

¹⁴Alkan (2002) and Alkan and Gale (2003) introduce related monotonicity conditions.

¹⁵Hatfield and Kojima (2010) introduce other notions of weak substitutability.

mechanism ϕ is denoted as $\phi_s(P_S)$. A mechanism ϕ satisfies *strategy-proofness* if no student can misreport her preferences and get a strictly more preferred contract. More formally, for every student *s* and preference profile P_S , there exists no preference P'_s such that $\phi_s(P'_s, P_{S \setminus \{s\}}) P_s \phi_s(P_S)$. For any property on matchings, a mechanism satisfies the property if, for every preference profile, the matching produced by the mechanism satisfies the property.

3. Achieving Policy Goals with Stable Outcomes

To achieve stable matchings with desirable properties, we use a generalization of the deferred-acceptance mechanism of Gale and Shapley (1962).

Student-Proposing Deferred Acceptance Mechanism (SPDA).

- **Step 1:** Each student *s* proposes a contract (s, d, c) to district *d* where *c* is her most preferred school. Let X_d^1 denote the set of contracts proposed to district *d*. District *d* tentatively accepts contracts in $Ch_d(X_d^1)$ and permanently rejects the rest. If there are no rejections, then stop and return $\cup_{d \in D} Ch_d(X_d^1)$ as the outcome.
- Step n (n > 1): Each student *s* whose contract was rejected in Step n 1 proposes a contract (s, d, c) to district *d* where *c* is her next preferred school. If there is no such school, then the student does not make any proposals. Let X_d^n denote the set of contracts that were tentatively accepted by district *d* in Step n - 1 and contracts that were proposed to district *d* in Step *n*. District *d* tentatively accepts contracts in $Ch_d(X_d^n)$ and permanently rejects the rest. If there are no rejections, then stop and return $\cup_{d \in D} Ch_d(X_d^n)$.

When district admissions rules have completions that satisfy substitutability and LAD, SPDA is stable and strategy-proof (Hatfield and Kominers, 2014). Therefore, when we analyze SPDA, we assume that students report their preferences truthfully.

We illustrate SPDA using the following example. We come back to this example later to study the effects of interdistrict school choice.

Example 1. Consider a problem with two school districts, d_1 and d_2 . District d_1 has school c_1 with capacity one and school c_2 with capacity two. District d_2 has school c_3 with capacity two. There are four students: students s_1 and s_2 are from district d_1 , whereas students s_3 and s_4 are from district d_2 . The initial matching is $\{(s_1, c_1), (s_2, c_2), (s_3, c_3), (s_4, c_3)\}$.

Given any set of contacts, district d_1 chooses students who have contracts with school c_1 first and then chooses from the remaining students who have contracts with school c_2 . For school c_1 , the district prioritizes students in the order $s_3 \succ s_4 \succ s_1 \succ s_2$ and chooses one applicant if there is any. For school c_2 , the district prioritizes students according to the order $s_1 \succ s_2 \succ s_3 \succ s_4$ and chooses as many applicants as possible without going over the school's capacity while ignoring the contracts of the students who have already been accepted at c_1 . Likewise, district d_2 prioritizes students according to the order $s_3 \succ s_4 \succ$ $s_1 \succ s_2$ and chooses as many applicants as possible without going over the capacity of school c_3 . These admissions rules are feasible and acceptant, and they have completions that satisfy substitutability and LAD.¹⁶ In addition, student preferences are given by the following table,

$\underline{P_{s_1}}$	$\underline{P_{s_2}}$	$\underline{P_{s_3}}$	P_{s_4}
c_1	c_3	c_1	c_2
c_2	c_1	c_2	c_1
c_3	c_2	c_3	c_3

which means that, for instance, student s_1 prefers c_1 to c_2 to c_3 .

In this problem, SPDA runs as follows. At the first step, student s_1 proposes to district d_1 with contract (s_1, c_1) , student s_2 proposes to district d_2 with contract (s_2, c_3) , student s_3 proposes to district d_1 with contract (s_3, c_1) , and student s_4 proposes to district d_1 with contract (s_4, c_2) . District d_1 first considers contracts associated with school c_1 , (s_1, c_1) and (s_3, c_1) , and tentatively accepts (s_3, c_1) while rejecting (s_1, c_1) because student s_3 has a higher priority than student s_1 at school c_2 . In this case, there is only one such contract, (s_4, c_2) , which is tentatively accepted. District d_2 considers contract (s_2, c_3) and tentatively accepted. District d_2 considers contract (s_2, c_3) and tentatively accepted. District d_2 considers contract (s_2, c_3) and tentatively accepts it. The tentative matching is $\{(s_2, c_3), (s_3, c_1), (s_4, c_2)\}$. Since there is a rejection, the algorithm proceeds to the next step.

At the second step, student s_1 proposes to district d_1 with contract (s_1, c_2) . District d_1 first considers contract (s_3, c_1) and tentatively accepts it. Then district d_1 considers contracts (s_1, c_2) and (s_4, c_2) and tentatively accepts them both. District d_2 does not have any new contracts, so tentatively accepts (s_2, c_3) . Since there is no rejection, the algorithm stops. The outcome of SPDA is $\{(s_1, c_2), (s_2, c_3), (s_3, c_1), (s_4, c_2)\}$.

In the rest of this section, we formalize three policy goals and characterize conditions under which SPDA satisfies them.

3.1. Individual Rationality. In our context, individual rationality requires that every student is matched with a weakly more preferred school than her initial school. As a result, SPDA does not necessarily satisfy individual rationality even though each student is either unmatched or matched with a school that is more preferred than being unmatched.

¹⁶In Appendix B.1, we provide a general class of admissions rules, including this one as a special case. We show that these admissions rules are feasible and acceptant, and they have completions that satisfy substitutability and LAD.

If individual rationality is violated so that some students prefer their initial schools to the outcome of SPDA, then there may be public opposition that harm interdistrict school choice efforts. For this reason, individual rationality is a desirable property for policymakers. The following condition proves to play a crucial role for achieving this property.

Definition 1. A district admissions rule Ch_d respects the initial matching if, for any student s whose initial school c is in district d and matching X that is feasible for students, $(s, d, c) \in X$ implies $(s, d, c) \in Ch_d(X)$.

When a district's admissions rule respects the initial matching, it has to admit those contracts associated with itself in which students apply to their initial schools. The following result shows that this is exactly the condition for SPDA to satisfy individual rationality.

Theorem 1. *SPDA satisfies individual rationality if, and only if, each district's admissions rule respects the initial matching.*

The intuition for the "if" part of this theorem is simple. When district admissions rules respect the initial matching, no student is matched with a school which is strictly less preferred than her initial school under SPDA because she is guaranteed to be accepted by that school if she applies to it. For the "only if" part of the theorem, we construct a specific student preference profile such that SPDA assigns one student a strictly less preferred school than her initial school whenever there exists one district with an admissions rule that does not respect the initial matching.

In the next example, we illustrate SPDA with district admissions rules that respect the initial matching.

Example 2. Consider the problem in Example 1. Recall that in this problem, the outcome of SPDA is $\{(s_1, c_2), (s_2, c_3), (s_3, c_1), (s_4, c_2)\}$. This matching is not individually rational because student s_1 prefers her initial school c_1 to school c_2 that she is matched with. This observation is consistent with Theorem 1 because the admissions rule of district d_1 does not respect the initial matching. In particular, $Ch_{d_1}(\{(s_1, c_1), (s_3, c_1)\}) = \{(s_3, c_1)\}$, so student s_1 is rejected from a matching that is feasible for students and includes the contract with her initial school.

Now modify the priority ranking of district d_1 at school c_1 so that $s_1 \succ s_2 \succ s_3 \succ s_4$ but, otherwise, keep the construction of the district admissions rules and student preferences the same as before. With this change, district admissions rules respect the initial matching because each student is accepted when she applies to the district with her initial school.¹⁷ In particular, the proposal of student s_1 to district d_1 with her initial school c_1 is

¹⁷In Appendix B.2, we construct a class of district admissions rules that includes this admissions rule as a special case. These admissions rules are feasible and acceptant, and have completions that satisfy substitutability and LAD. Furthermore, they also respect the initial matching.

always accepted. With this modification, it is easy to check that the outcome of SPDA is $\{(s_1, c_1), (s_2, c_3), (s_3, c_2), (s_4, c_2)\}$. This matching satisfies individual rationality.

In some school districts, each student gets a priority at her neighborhood school, as in this example. In the absence of other types of priorities, neighborhood priority guarantees that SPDA satisfies individual rationality.

3.2. Balanced Exchange. For an interdistrict school choice program, maintaining a balance of students incoming from and outgoing to other districts is important. To formalize this idea, we say that a mechanism satisfies the *balanced-exchange* policy if the number of students that a district gets from the other districts and the number of students that the district sends to the others are the same for every district and for every profile of student preferences. Since district choice rules are acceptant, every student is matched with a school under SPDA. Therefore, for SPDA, this policy is equivalent to the requirement that the number of students assigned to a district must be equal to the number of students from that district.

The balanced-exchange policy is important because the funding that a district gets depends on the number of students it serves. Therefore, an interdistrict school choice program may not be sustainable if SPDA does not satisfy the balanced-exchange policy. For achieving this policy goal, the following condition on admissions rules proves important.

Definition 2. A matching X is **rationed** if, for every district d, it does not assign strictly more students to the district than the number of students whose home district is d. A district admissions rule is **rationed** if it chooses a rationed matching from any matching that is feasible for students.

When a district admissions rule is rationed, then the district does not accept strictly more students than the number of students from the district at any matching that is feasible for students. The result below establishes that this property is exactly the condition to guarantee that SPDA satisfies the balanced-exchange policy.

Theorem 2. *SPDA satisfies the balanced-exchange policy if, and only if, each district's admissions rule is rationed.*

To obtain the intuition for this result, consider a student. Acceptance requires that a district can reject all contracts of this student only when the number of students assigned to the district is at least as large as the number of students from that district. As a result, all students are guaranteed to be matched with some school district. In addition, when district admissions rules are rationed, a district cannot accept more students than the number of students from the district. These two facts together imply that the number of students assigned to a district in SPDA is equal to the number of students from that district. Therefore, SPDA satisfies the balanced-exchange policy when each district's admissions rule is

rationed. Conversely, when there exists one district with an admissions rule that fails to be rationed, then we can construct student preferences such that this district is matched with strictly more students than the number of students from the district in SPDA, which means that the outcome does not satisfy the balanced-exchange policy.

Now we illustrate SPDA when district admissions rules are rationed.

Example 3. Consider the problem in Example 1. Recall that in this problem, the SPDA outcome is $\{(s_1, c_2), (s_2, c_3), (s_3, c_1), (s_4, c_2)\}$. Since there are three students matched with district d_1 and there are only two students from that district, SPDA does not satisfy the balanced-exchange policy. This is consistent with Theorem 2 because the admissions rule of district d_1 is not rationed. In particular, $Ch_{d_1}(\{(s_1, c_2), (s_3, c_1), (s_4, c_2)\}) = \{(s_1, c_2), (s_3, c_1), (s_4, c_2)\}$, so district d_1 accepts more students than the number of students from there given a matching that is feasible for students.

Suppose that we modify the admissions rule of district d_1 as follows. If the district chooses a contract associated with school c_1 , then at most one contract associated with school c_2 is chosen. Therefore, the district never chooses more than two contracts, which is the number of students from there. Therefore, the updated admissions rule is rationed.¹⁸ With this change, it is easy to check that the SPDA outcome is $\{(s_1, c_2), (s_2, c_3), (s_3, c_1), (s_4, c_3)\}$, which satisfies the balanced-exchange policy.

An implication of Theorems 1 and 2 is that SPDA is guaranteed to satisfy individual rationality and the balanced-exchange policy if, and only if, each district's admissions rule respects the initial matching and is rationed.

3.3. Diversity. The third policy goal we consider is that of diversity. More specifically, we are interested in how to ensure that there is enough diversity across districts so that the student composition in terms of demographics does not vary too much from district to district.

We are mainly motivated by a program that is used in the state of Minnesota. State law in Minnesota identifies racially isolated (relative to one of their neighbors) school districts and requires them to be in the *Achievement and Integration (AI) Program*. The goal is to increase the racial parity between neighboring school districts. We first introduce a diversity policy in the spirit of this program: Given a constant $\alpha \in [0, 1]$, we say that a mechanism satisfies the α -diversity policy if for all preferences, districts d and d', and type t, the difference between the ratios of type-t students in districts d and d' is not more than α . We interpret α to be the maximum ratio difference tolerated under the diversity policy; for instance, $\alpha = 0.2$ for Minnesota.

¹⁸In Appendix B.3, we construct a class of rationed district admissions rules that includes this admissions rule as a special case. These admissions rules are feasible and acceptant, and they have completions that satisfy substitutability and LAD.

We study admissions rules such that SPDA satisfies the α -diversity policy when there is interdistrict school choice. Since this policy restricts the number of students across districts, a natural starting point is to have type-specific ceilings at the district level. However, it turns out that type-specific ceilings at the district level may result in district admissions rules resulting in no stable matchings (see Theorem 9 in Appendix A.2).

Since there is an incompatibility between district-level type-specific ceilings and the existence of a stable matching, we impose type-specific ceilings at the school level as follows.

Definition 3. A district admissions rule Ch_d has a school-level type-specific ceiling of q_c^t at school c for type-t students if the number of type-t students admitted cannot exceed this ceiling. More formally, for any matching X that is feasible for students,

$$|\{x \in Ch_d(X) | \tau(s(x)) = t, c(x) = c\}| \le q_c^t$$

Note that district admissions rules typically violate once school-level type-specific ceilings are imposed. This is because a student can be rejected from a set that is feasible for students even when the number of applicants to each school is weakly smaller than its capacity and the number of applicants to the district is weakly smaller than the number of students from that district. Given this, we define a weaker version of the acceptance assumption as follows.

Definition 4. A district admissions rule Ch_d that has school-level type-specific ceilings is **weakly** *acceptant* if, for any contract x associated with a type-t student and district d and matching X that is feasible for students, if x is rejected from X, then at $Ch_d(X)$,

- the number of students assigned to school c(x) is equal to $q_{c(x)}$, or
- the number of students assigned to district d is at least k_d , or
- the number of type-t students assigned to school c(x) is at least q_c^t .

In other words, a student can be rejected from a set that is feasible for students only when one of these three conditions is satisfied.

In SPDA, a student may be left unassigned due to school-level type-specific ceilings even when district admissions rules are weakly acceptant. To make sure that every student is matched, we make the following assumption.

Definition 5. District admissions rules $(Ch_d)_{d \in D}$ accommodate unmatched students if for any student *s* and feasible matching *X* in which student *s* is unmatched, there exists $x = (s, d, c) \in \mathcal{X}$ such that $x \in Ch_d(X \cup \{x\})$.

When district admissions rules accommodate unmatched students, for any feasible matching in which a student is unmatched, there exists a school such that the district associated with the school would admit that student if she applies to that school. For example, when admissions rules respect the initial matching, they also accommodate unmatched

students because an unmatched student's application to her initial school is always accepted. Lemma 2 in Appendix D shows that when district admissions rules accommodate unmatched students, every student is matched to a school in SPDA.

In general, accommodation of unmatched students may be in conflict with type-specific ceilings because there may not be enough space for a student type when ceilings are small for this type. To avoid this, we assume that type-specific ceilings are high enough so that $(Ch_d)_{d\in\mathcal{D}}$ accommodate unmatched students.¹⁹

Our assumptions on district admissions rules allow us to control the distribution of the SPDA outcome. In particular, the SPDA outcome satisfies the following four conditions. We call any matching X satisfying those conditions **legitimate**: (i) $\sum_t \xi_d^t(X) = k_d$ for all $d \in \mathcal{D}$, (ii) $\sum_{c \in \mathcal{C}} \xi_c^t(X) = k^t$ for all $t \in \mathcal{T}$, (iii) $\sum_{t \in \mathcal{T}} \xi_c^t(X) \leq q_c$ for all $c \in \mathcal{C}$, and (iv) $\xi_c^t(X) \leq q_c^t$ for all $t \in \mathcal{T}$ and $d \in \mathcal{D}$.

In this framework, type-*t* ceilings of schools in district *d* may result in a floor of another type *t'* in this district in the sense that the number of type-*t'* students in the district should be at least a certain number. Moreover, this may further impose a ceiling for type *t'* in another district *d'*. To see this, suppose, for example, that (i) there are two districts *d* and *d'*, (ii) in each district, there is one school and 100 students, (iii) 100 students are of type *t* and 100 students are of another type *t'*, and (iv) each school has a type-*t* ceiling of 60 and a type-*t'* ceiling of 70. In a legitimate matching, each district needs to have at least 40 type-*t'* students (because, otherwise, the number of type-*t* students in that district would have to be more than 60). Moreover, this would mean that there cannot be more than 60 type-*t'* students in any district (because, otherwise, there would need to be more than 40 type-*t'* students in the other district, contradicting the floor we just calculated). Hence, in this example, in effect we have a floor of 40 and a (further restricted) ceiling of 60 for type-*t'* students for each district.

Faced with this complication, our approach is to find the tightest lower and upper bounds induced by these constraints. For this purpose, a certain optimization problem proves useful. More specifically, consider a linear-programming problem where for each type *t* and district *d*, we seek the minimum and maximum values of $\sum_{c:d(c)=d} y_c^t$ subject to (i) $\sum_{t'\in\mathcal{T}}\sum_{c:d(c)=d'} y_c^{t'} = k_{d'}$ for all $d' \in \mathcal{D}$, (ii) $\sum_{c\in\mathcal{C}} y_c^{t'} = k^{t'}$ for all $t' \in \mathcal{T}$, (iii) $\sum_{t'\in\mathcal{T}} y_c^{t'} \leq q_c$ for all $c \in \mathcal{C}$, and (vi) $y_c^{t'} \leq q_c^{t'}$ for all $t' \in \mathcal{T}$ and $d' \in \mathcal{D}$. Let \hat{p}_d^t and \hat{q}_d^t be the solutions to the minimization and maximization problems, respectively.

¹⁹For instance, ignoring integer problems, $q_d^t \ge k_d \frac{k^t}{\sum_{t' \in \mathcal{T}} k^{t'}}$ for all t, d, would make ceilings compatible with this property as it would be possible to assign the same percentage of students of each type to all districts.

Both of these optimization problems belong to a special class of linear-programming problems called a minimum-cost flow problem, and many computationally efficient algorithms to solve it are known in the literature.²⁰ A straightforward but important observation is that \hat{p}_d^t (resp. \hat{q}_d^t) is exactly the lowest (resp. highest) number of type-*t* students who can be matched to district *d* in a legitimate matching (Lemma 3 in Appendix D). Given this observation, we call \hat{p}_d^t the *implied floor* and \hat{q}_d^t the *implied ceiling*.

Now we are ready to state the main result of this section.

Theorem 3. Suppose that each district admissions rule has school-level type-specific ceilings and is rationed and weakly acceptant. Moreover, suppose that the district admissions rules accommodate unmatched students. SPDA satisfies the α -diversity policy if, and only if, $\hat{q}_d^t/k_d - \hat{p}_{d'}^t/k_{d'} \leq \alpha$ for every type t and districts d, d' such that $d \neq d'$.

The proof of this theorem, given in Appendix D, is based on a number of steps. First, as mentioned above, we note that \hat{p}_d^t and \hat{q}_d^t are the lower and upper bounds, respectively, of the numbers of type-*t* students who can be matched in district *d* in any legitimate matching. This observation immediately establishes the first part of the theorem. Then, we further establish that the implied floors and ceilings are not arbitrary lower and upper bounds but can be achieved simultaneously in the sense that, for any pair of districts *d* and *d'* with $d \neq d'$, there exists a legitimate matching that assigns exactly \hat{q}_d^t type-*t* students to district *d* and exactly $\hat{p}_{d'}^t$ type-*t* students to district *d'* (Lemma 4). In other words, we establish that the implied ceiling and floor are achieved in two different districts, and they are achieved at *one* matching at the same time. We complete the proof of the theorem by constructing preferences such that the outcome of SPDA achieves these bounds. In Appendix C, we provide an example that illustrates Theorem 3.

The analysis in this section characterizes conditions under which different policy goals are achieved under SPDA. One of the facts worth mentioning in this context is that achieving multiple policies can be overly demanding. To see this point, we note that individual rationality and α -diversity policy are often incompatible with one another. For example, consider a problem such that each student's most preferred school is her initial school and a constant α such that the initial matching does not satisfy the α -diversity policy. Indeed,

²⁰To see that our problem is a minimum-cost flow problem, note that we can take $(k_d)_{d\in\mathcal{D}}$ as the "supply," $(k^t)_{t\in\mathcal{T}}$ as the "demand," $(q_d^t)_{d\in\mathcal{D},t\in\mathcal{T}}$ as the "arc capacity bounds," and the objective functions for \hat{p}_d^t and \hat{q}_d^t to be min y_d^t and min $-y_d^t$, respectively. These problems have an "integrality property" so that if the supply, demand, and bounds are integers, then all the solutions are integers as well. As already mentioned, many algorithms have been proposed to solve different objective functions for these problems. For instance, the capacity scaling algorithm of Edmonds and Karp (1972) gives the solutions in polynomial time. For more information, see Chapter 10 of Ahuja (2017). We are grateful to Fatma Kilinc-Karzan for helpful discussions.

in this case, no mechanism can simultaneously satisfy individual rationality and the α diversity policy because the initial matching is the unique individually rational matching.

4. Achieving Policy Goals with Efficient Outcomes

In this section, we turn our focus to efficiency. More specifically, we study the existence of a mechanism that satisfies a given policy goal on the distribution of agents, constrained efficiency, strategy-proofness, and individual rationality. We first consider a policy goal with type-specific ceilings at the district level. In this setting, we establish an impossibility result.

Theorem 4. There exist a problem and ceilings $(q_d^t)_{t \in \mathcal{T}, d \in \mathcal{D}}$ for which no mechanism satisfies the policy goal $\Xi \equiv \{\xi | q_d^t \ge \xi_d^t \text{ for all } d \text{ and } t, q_c \ge \sum_t \xi_c^t \text{ for all } c\}$, constrained efficiency, individual rationality, and strategy-proofness.

We show this result using the following example.

Example 4. Consider the following problem with districts d_1 and d_2 . District d_1 has schools c_1 , c_2 , and c_3 and district d_2 has schools c_4 , c_5 , and c_6 . All schools have a capacity of one. There are six students: students s_1 and s_4 have type t_1 , students s_2 and s_5 have type t_2 , and students s_3 and s_6 have type t_3 . Both districts have a ceiling of one for types t_1 and t_2 : $q_{d_1}^{t_1} = q_{d_1}^{t_2} = 1$ and $q_{d_2}^{t_1} = q_{d_2}^{t_2} = 1$. Initially, student s_i is matched with school c_i , for $i = 1, \ldots, 6$. Student preferences are as follows:

$\underline{s_1}$	$\underline{s_2}$	$\underline{s_3}$	$\underline{s_4}$	$\underline{s_5}$	$\underline{s_6}$
c_6	c_6	c_5	c_3	c_3	c_1
c_1	c_2	c_4	c_4	c_5	c_2
÷	÷	c_3	÷	÷	c_6
		÷			÷

where the dots in the table mean that the corresponding parts of the preferences are arbitrary.

In this example, there are two matchings that satisfy the policy goal Ξ , constrained efficiency, and individual rationality:

$$X = \{(s_1, c_6), (s_2, c_2), (s_3, c_4), (s_4, c_3), (s_5, c_5), (s_6, c_1)\}, \text{ and}$$
$$X' = \{(s_1, c_1), (s_2, c_6), (s_3, c_5), (s_4, c_4), (s_5, c_3), (s_6, c_2)\}.$$

If a mechanism satisfies the desired properties, then its outcome at the above student preference profile must be either matching X or X'.

Consider the case where the mechanism produces matching X at the above student preference profile. Suppose student s_3 misreports her preference by ranking c_5 first and c_3 second (while the ranking of other schools is arbitrary). Under the new report, the mechanism produces matching X' because it is the only matching that satisfies the policy goal Ξ , constrained-efficiency, and individual rationality. Since student s_3 strictly prefers her school in X' to her school in X, she has a profitable deviation.

Similarly, consider the case where the mechanism produces matching X' at the above student preference profile. Suppose student s_6 misreports her preference by ranking c_1 first and c_6 second (while the ranking of other schools arbitrary). In this case, the mechanism produces matching X because it is the only matching that satisfies the policy goal Ξ , constrained-efficiency, and individual rationality. Since student s_6 strictly prefers her school in X to her school in X', she has a profitable deviation.

In both cases, there exists a student who benefits from misreporting, so the desired conclusion follows. $\hfill \Box$

This example also shows that there is no mechanism that satisfies the α -diversity policy goal for $\alpha = 0$ introduced in Section 3.3, constrained efficiency, individual rationality, and strategy-proofness. Consequently, without any assumptions, a policy goal may not be implemented with the desirable properties. To establish a positive result, we consider distributional policy goals that satisfy the following notion of discrete convexity, which is studied in the mathematics and operations research literatures (Murota, 2003).

Definition 6. Let $\chi_{c,t}$ denote the distribution where there is one type-t student at school c and there are no other students. A set of distributions Ξ is **M-convex** if whenever $\xi, \tilde{\xi} \in \Xi$ and $\xi_c^t > \tilde{\xi}_c^t$ for some school c and type t then there exist school c' and type t' with $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ such that $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi$ and $\tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi$.²¹

To illustrate this concept, suppose that a set of distributions Ξ is M-convex. Consider two distributions ξ and $\tilde{\xi}$ in this set such that there are more type-*t* students in school *c* at ξ than at $\tilde{\xi}$. Then there exist school *c'* and type *t'* such that there are more type-*t'* students in school *c'* at $\tilde{\xi}$ than ξ with the following two properties. First, removing one type-*t* student from school *c* and adding one type-*t'* student to school *c'* in ξ produces a distribution in Ξ . Second, removing one type-*t'* student from school *c'* and adding one type-*t* student to school *c* in $\tilde{\xi}$ gives a distribution in Ξ (see Figure 2). Intuitively, from each of these two distributions we can move closer to the other distribution in an incremental manner within Ξ , a property analogous to the standard convexity notion but adapted to a discrete setting. We illustrate this concept with the following example.

²¹The letter M in the term M-convex set comes from the word *matroid*, a closely related and well-studied concept in discrete mathematics.



FIGURE 2. Illustration of M-convexity

Example 5. Consider the problem and the set of distributions Ξ defined in Example 4. We show that Ξ is not M-convex. Recall matchings X and X' in that example. By construction, both X and X' satisfy the policy goal Ξ . Furthermore, $\xi_{c_3}^{t_1}(X) = 1 > 0 = \xi_{c_3}^{t_1}(X')$ because (i) school c_3 is matched with student s_4 at X, whose type is t_1 , while (ii) school c_3 is matched with student s_4 at X, whose type is t_1 , while (ii) school c_3 is matched with student s_5 at X', whose type is $t_2 \neq t_1$. If the set of distributions Ξ is M-convex, there exist a school c and a type t such that $\xi_c^t(X) < \xi_c^t(X')$ and $\xi(X) - \chi_{c_3,t_1} + \chi_{c,t}$ is in Ξ . Because each school's capacity is one, and at matching X all schools have filled their capacities, this means that the only candidate for (c, t) satisfying the above condition is such that $c = c_3$. But the only nonzero $\xi_{c_3}^t(X')$ is for $t = t_2$ (corresponding to s_5 matched with c_3 at X'), and $\xi(X) - \chi_{c_3,t_1} + \chi_{c_3,t_2}$ does not satisfy the policy goal because district d_1 's ceiling for type t_2 is violated (note $\xi_{c_2}^{t_2}(X) = 1$ because student s_2 is matched with c_2 at X.)

The above argument implies that $\Xi \cap \Xi^0$ is not M-convex either. To see this, note that both $\xi(X)$ and $\xi(X')$ are in $\Xi \cap \Xi^0$ because all students are matched. Because we have shown that no distribution of the form $\xi(X) - \chi_{c_3,t_1} + \chi_{c,t}$ is in Ξ , by set inclusion relation $\Xi \cap \Xi^0 \subseteq \Xi$, there is no distribution of the form $\xi(X) - \chi_{c_3,t_1} + \chi_{c,t}$ in $\Xi \cap \Xi^0$ either. \Box

Now we introduce an algorithm that achieves the desirable properties whenever the policy goal is M-convex. To do this, we first create a hypothetical matching problem. On one side of the market, there are school-type pairs (c,t) where $c \in C$ and $t \in T$. On the other side, there are students from the original problem, S. Given any student $s \in S$ and a preference order P_s of s in the original problem, define preference order \tilde{P}_s over school-type pairs in the hypothetical problem as follows: letting t be the type of student s and c_0 be her initial school in the original problem, $(s,c) P_s(s,c') \iff (c,t) \tilde{P}_s(c',t)$ for any

 $c, c' \in C$, and $(c_0, t) \tilde{P}_s(c, t')$ for any $c \in C$ and $t' \in T$ such that $t' \neq t$. That is, \tilde{P}_s is a preference order over school-type pairs that ranks the school-type pairs in which the type is t in the same order as in P_s , while finding all school-type pairs specifying a different type as less preferred than the pair corresponding to her initial school. Furthermore, let (c_0, t) be the initial school for s in the hypothetical problem.

Next we define a priority ordering of students that school-type pairs use to rank students. For school-type pair (c, t), students initially matched with (c, t) have the highest priority, and then all other students have the second highest priority. This gives us two priority classes for students. Then, ties are broken according to a master priority list that every school-type pair uses.

We say that a type-*t* student *s* with initial matching school-type pair (c, t) is *permissible* to school-type pair (c', t') at matching *X* if $\xi(X) + \chi_{c',t'} - \chi_{c,t}$ is in Ξ . Note that a type-*t* student with initial matching school-type pair (c, t) is always permissible to pair (c, t) at matching *X* whenever $\xi(X)$ is in Ξ .

The following is a generalization of Gale's top trading cycles algorithm (Shapley and Scarf, 1974), building on its recent extension by Suzuki et al. (2017).

Top Trading Cycles Algorithm (TTC). Consider a hypothetical problem.

- **Step 1:** Let $X^1 \equiv \tilde{X}$. Each school-type pair points to the permissible student at matching X^1 with the highest priority. If there exists no such student, remove the school-type pair from the market. Each student *s* points to the highest ranked remaining school-type pair with respect to \tilde{P}_s . Identify and execute cycles. Any student who is part of an executed cycle is assigned the school-type pair she is pointing to and is removed from the market.
- Step n (n > 1): Let X^n denote the matching consisting of matchings assigned in the previous steps and initial matchings for all students who have not been processed in the previous steps. Each remaining school-type pair points to the unassigned student who is permissible at matching X^n with the highest priority. If there exists no such student, remove the school-type pair from the market. Each unassigned student *s* points to the highest ranked remaining school-type pair with respect to \tilde{P}_s . Identify and execute cycles. Any student who is part of an executed cycle is matched with the school-type pair she is pointing to and is removed from the market.

This algorithm terminates in the first step such that no student remains to be processed. The TTC outcome is defined as the matching induced by the outcome of the hypothetical problem at this step. Note that the definition of permissibility and, hence, the definition of TTC, depend on the policy goal. Nevertheless, we do not explicitly state the policy goal under consideration when it is clear from the context.

The main result of this section is as follows.

Theorem 5. Suppose that the initial matching satisfies the policy goal Ξ . If $\Xi \cap \Xi^0$ is M-convex, then TTC satisfies the policy goal Ξ , constrained efficiency, individual rationality, and strategy-proofness.

The assumption that the initial matching satisfies the policy goal is necessary for the result: Consider student preferences such that each student's highest-ranked school is her initial school. Then the initial matching is the unique individually rational matching. Therefore, if there exists a mechanism with the desired properties, then the outcome at this preference profile has to be the initial matching. Hence, we need the assumption that the initial matching satisfies the policy goal to have such a mechanism.

To see one of the implications of this theorem, suppose that the policy goal Ξ is such that no school is matched with more students than its capacity. In that case, if Ξ is M-convex, then TTC satisfies the desirable properties.

Corollary 1. Suppose that the policy goal Ξ is such that for every $\xi \in \Xi$ and $c \in C$, $\sum_t \xi_c^t \leq q_c$. Furthermore, suppose that the initial matching satisfies Ξ . If Ξ is M-convex, then TTC satisfies the policy goal Ξ , constrained efficiency, individual rationality, and strategy-proofness.

In the proof of this corollary, we show that when Ξ is M-convex and no distribution in Ξ assigns more students to a school than its capacity, then $\Xi \cap \Xi^0$ is also M-convex. Therefore, the corollary follows directly from Theorem 5.

Next we illustrate TTC with an example.

Example 6. Consider a problem with two school districts, d_1 and d_2 . District d_1 has school c_1 with capacity three and school c_2 with capacity two. District d_2 has school c_3 with capacity two and school c_4 with capacity one. There are seven students: students s_1 , s_2 , s_3 , and s_4 are from district d_1 and have type t_1 , whereas students s_5 , s_6 , and s_7 are from district d_2 and have type t_2 . The initial matching is $\{(s_1, c_1), (s_2, c_1), (s_3, c_2), (s_4, c_2), (s_5, c_3), (s_6, c_3), (s_7, c_4)\}$. Student preferences are as follows.

$\underline{P_{s_1}}$	$\underline{P_{s_2}}$	P_{s_3}	$\underline{P_{s_4}}$	P_{s_5}	$\underline{P_{s_6}}$	P_{s_7}
c_2	c_3	c_4	c_2	c_1	c_4	c_2
c_3	c_1	c_2	c_3	c_2	c_1	c_3
:	÷	÷	c_1	c_3	c_3	c_1
			c_4	c_4	c_2	c_4

In addition to the school capacities, there is only one additional constraint that school c_1 cannot have more than one type- t_2 student. As we show in the proof of Corollary 2, the

set of distributions that satisfy this policy goal and the requirement that every student is matched is an M-convex set. Therefore, by Theorem 5, TTC satisfies constrained efficiency, individual rationality, strategy-proofness, and the policy goal.

To run TTC, we use a master priority list. Suppose that the master priority list ranks students as follows: $s_1 \succ s_2 \succ s_3 \succ s_4 \succ s_5 \succ s_6 \succ s_7$.

At Step 1 of TTC, there are eight school-type pairs. Consider (c_1, t_1) . Initially, students s_1 and s_2 are matched with it, so they are both permissible to this pair. We use the master priority list to rank them, so s_1 gets the highest priority at (c_1, t_1) . Therefore, (c_1, t_1) points to s_1 . Now consider (c_1, t_2) . Initially, it does not have any students because there is no type- t_2 student assigned to c_1 in the original problem. Furthermore, s_1 is permissible to (c_1, t_2) because she can be removed from (c_1, t_1) and a type- t_2 student can be assigned to (c_1, t_2) without violating the school capacities or the policy goal. Therefore, (c_1, t_2) points to s_1 as well, who gets a higher priority than the other permissible students because of the master priority list. The rest of the pairs also point to the highest-priority permissible students. Each student points to the highest ranked school-type pair of the same type as shown in Figure 3A. There is only one cycle: $s_7 \rightarrow (c_2, t_2) \rightarrow s_3 \rightarrow (c_4, t_1) \rightarrow s_7$. Therefore, s_7 is matched with (c_2, t_2) and s_3 is matched with (c_4, t_1) .

At Step 2, there are six remaining school-type pairs: There are no permissible students for (c_4, t_1) and (c_4, t_2) because c_4 has a capacity of one and it is already assigned to s_3 . Each remaining school-type pair points to the highest-ranked remaining permissible student. Each student points to the highest-ranked remaining school-type pair (see Figure 3B). There is only one cycle: $s_4 \rightarrow (c_2, t_1) \rightarrow s_4$. Hence, s_4 is assigned to (c_2, t_1) .

The algorithm ends in five steps. Steps 3 and 4 are also shown in Figure 3. In Step 5, s_2 points to (c_1, t_1) , which points back to the student. The outcome of the algorithm is

$$\{(s_1, c_3), (s_2, c_1), (s_3, c_4), (s_4, c_2), (s_5, c_1), (s_6, c_3), (s_7, c_2)\}.$$

It can be easily seen that the distribution associated with this matching satisfies the policy goal because no school has more students than its capacity and c_1 has only one type- t_2 student.

Sometimes it may be more convenient to describe a policy goal using a real-valued function rather than a set of distributions. The interpretation is that the policy function measures how satisfactory the distribution is in terms of the policy goal. To formalize this alternative approach let $f : \mathbb{Z}_{+}^{|\mathcal{C}| \times |\mathcal{T}|} \to \mathbb{R}$ be a function on distributions such that $f(\xi) \ge f(\xi')$ means that distribution ξ satisfies the policy at least as well as distribution ξ' . Let $\lambda \in \mathbb{R}$ be a constant. Consider the following (f, λ) -policy: $\Xi(f, \lambda) \equiv \{\xi \in \mathbb{Z}_{+}^{|\mathcal{C}| \times |\mathcal{T}|} | f(\xi) \ge \lambda\}$. Note that the initial matching \tilde{X} satisfies the (f, λ) -policy if, and only if, $f(\xi(\tilde{X})) \ge \lambda$.



FIGURE 3. The first four steps of TTC. In each step, there is only one cycle, which is represented by the dashed lines.

We introduce the following condition on functions, which plays a crucial role in the M-convexity of the (f, λ) -policy.

Definition 7. A function f is pseudo M-concave, if for every distinct $\xi, \tilde{\xi} \in \Xi_0$, there exist (c, t) and (c', t') with $\xi_c^t > \tilde{\xi}_c^t$ and $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ such that

$$\min\{f(\xi - \chi_{c,t} + \chi_{c',t'}), f(\hat{\xi} + \chi_{c,t} - \chi_{c',t'})\} \ge \min\{f(\xi), f(\hat{\xi})\}.$$

This is a notion of concavity for functions on a discrete domain. Lemma 1 shows that pseudo M-concavity characterizes when upper contour sets are M-convex. It is stronger than quasi M-concavity but not logically related to the M-concavity studied in the discrete mathematics literature (Murota, 2003).

Lemma 1. $\Xi(f, \lambda) \cap \Xi^0$ is *M*-convex for every λ if, and only if, f is pseudo *M*-concave.

Therefore, we get the following result:

Theorem 6. If f is pseudo M-concave and λ is such that $f(\xi(\tilde{X})) \geq \lambda$, then TTC satisfies the (f, λ) -policy, constrained efficiency, individual rationality, and strategy-proofness.

To see why this theorem holds, recall that by Lemma 1, $\Xi(f, \lambda) \cap \Xi^0$ is M-convex. Furthermore, the initial matching satisfies the (f, λ) -policy. Therefore, the result follows from Theorem 5.

Before we consider specific policy goals, we show that the set-based approach in Theorem 5 and the function-based approach in Theorem 6 are equivalent. For that purpose, note first that Lemma 1 already shows that the (f, λ) -policy for a pseudo M-concave function f yields an M-convex policy set $\Xi(f, \lambda) \cap \Xi^0$. We establish a sense in which a converse result holds.

Theorem 7. Suppose that Ξ is a set of distributions. If $\Xi \cap \Xi^0$ is M-convex, then there exist a pseudo M-concave function f and a constant $\lambda \in \mathbb{R}$ such that $\Xi(f, \lambda) \cap \Xi^0 = \Xi \cap \Xi^0$.

Now that we have established general results based on M-convexity of the policy set or pseudo M-concavity of the policy function, we proceed to apply them to a variety of situations. To begin, consider the set Ξ of distributions of all feasible matchings. In other words, consider a situation in which no policy goal is imposed other than $\sum_t \xi_c^t \leq q_c$ for each c. Then it is rather straightforward to show that the set $\Xi \cap \Xi^0$ is an M-convex set. This implies that when there is no policy goal, TTC is efficient, individually rational, and strategy-proof, a standard result in the literature (Abdulkadiroğlu and Sönmez, 2003).

Next we apply Theorems 5 and 6 to a variety of policy goals. These results turn out to be applicable to many specific cases, as a wide variety of policy goals induce distributions that satisfy M-convexity or can be expressed by policy functions that are pseudo M-concave. To be more specific, first suppose that the policy goal Ξ sets type-specific floors and ceilings at each school, i.e., $\Xi \equiv \{\xi | q_c^t \ge \xi_c^t \ge p_c^t \text{ for all } c \text{ and } t, q_c \ge \sum_t \xi_c^t \text{ for all } c\}$ where q_c^t is the ceiling and p_c^t is the floor for type t at school c. Therefore, for each school, the number of students of a given type must be within the ceiling and floor of this type at the school. We call a policy goal Ξ of this form a *school-level diversity policy* and show that $\Xi \cap \Xi^0$ is an M-convex set. This finding, together with Theorem 5, implies the following positive result.

Corollary 2. Suppose that the initial matching satisfies a school-level diversity policy. Then TTC satisfies the school-level diversity policy, constrained efficiency, individual rationality, and strategy-proofness.

We note a sharp contrast between this result and Theorem 4. The latter result demonstrates that no mechanism is guaranteed to satisfy the policy goal and other desiderata such as constrained efficiency, individual rationality, and strategy-proofness if the floors or ceilings are imposed at the district level. Corollary 2, in contrast, shows that a mechanism with the desirable properties exists if the floors and ceilings are imposed at the school level. Taken together, these results inform policy makers about what kinds of diversity policies are compatible with the other desiderata.

One possible shortcoming of Corollary 2 is that the result holds under the assumption that the initial matching satisfies the school-level diversity policy. This may be undesirable given that often diversity policies are implemented because schools or districts are regarded as insufficiently diverse, as in the case of the diversity law in Minnesota. In such a setting, a potential diversity requirement can be that the diversity should not decrease as a result of interdistrict school choice according to a diversity measure f. Such a consideration can be formally described as the $(f, \xi(\tilde{X}))$ -policy, $\Xi(f, \xi(\tilde{X}))$. The next corollary establishes a positive result for a $\Xi(f, \xi(\tilde{X}))$ -policy where the diversity is measured via the "Manhattan distance" to an ideal point.

Corollary 3. Let $\hat{\xi} \in \Xi_0$ be an ideal distribution and $f(\xi) \equiv -\sum_{c,t} |\xi_c^t - \hat{\xi}_c^t|$ be the policy function. Then TTC satisfies $(f, \xi(\tilde{X}))$ -policy, constrained efficiency, individual rationality, and strategy-proofness.

Note that the initial matching \tilde{X} always satisfies $(f, \xi(\tilde{X}))$ -policy. Furthermore, we show that the policy function f is pseudo M-concave. Therefore, this corollary follows from Theorem 6. More generally, when the diversity is measured by a pseudo M-concave function, then the TTC outcome is as diverse as the initial matching. Furthermore, TTC also satisfies the other desirable properties.

Next, we study the balanced-exchange policy introduced in Section 3.2. We establish that the balanced-exchange policy is represented by a distribution that satisfies Mconvexity. This implies the following result.

Corollary 4. TTC satisfies the balanced-exchange policy, constrained efficiency, individual rationality, and strategy-proofness.

One of the advantages of our approach is that M-convexity of a set and pseudo Mconcavity of a function are so general that a wide variety of policy goals satisfy them, and that it is likely to be applicable for policy goals that one may encounter in the future. To highlight this point, we consider imposing the diversity and balanced-exchange policies at the same time. More specifically, define a set of distributions $\Xi \equiv \{\xi | q_c^t \ge \xi_c^t \ge p_c^t \text{ for all } c \text{ and } t, q_c \ge \sum_t \xi_c^t \text{ for all } c, \text{ and } \sum_t \sum_{c:d(c)=d} \xi_c^t = k_d \text{ for all } d\}$ and call it the *combination of balanced-exchange and school-level diversity policies*. This is the set of distributions that satisfy both the school-level floors and ceilings and the balanced-exchange requirement. We can establish this set is M-convex, implying the following result.

Corollary 5. Suppose that the initial matching satisfies the combination of balanced exchange and school-level diversity policies. Then TTC satisfies the combination of balanced exchange and school-level diversity policies, constrained efficiency, individual rationality, and strategy-proofness.

In the proof, we show that the combination of balanced exchange and school-level diversity policies satisfies M-convexity. In general, the intersection of two M-convex sets need not be M-convex.²² Therefore, the M-convexity of the combination of balanced exchange and school-level diversity policies does not follow from the M-convexity of the balanced exchange policy and the M-convexity of school-level diversity policy.

5. Conclusion

Despite increasing interest in interdistrict school choice in the US, the scope of matching theory has been limited to intradistrict choice. In this paper, we proposed a new frame-work to study interdistrict school choice that allows for interdistrict admissions, both from stability and efficiency perspectives. For stable mechanisms, we characterized conditions on district admissions rules that achieve a variety of important policy goals, such as student diversity across districts. For efficient mechanisms, we showed that certain types of diversity policies are incompatible with desirable properties such as strategy-proofness, while alternative forms of diversity policies can be achieved by a strategy-proof mechanism: a variation of the top trading cycles algorithm. Overall, our analysis suggests that interdistrict school choice can help achieve desirable policy goals such as student diversity, but only with an appropriate design of constraints, admissions rules, and placement mechanisms.

We regard this paper as a first step toward formal analysis of interdistrict school choice based on tools of market design. As such, we envision a variety of directions for future research. For example, it may be interesting to study cases in which the conditions for our results are violated. Although we already know the policy goals are not guaranteed to be satisfied for our stability results (our results provide necessary and sufficient conditions), the seriousness of the failure of the policy goals studied in the present paper is an open question. Quantitative measures or an approximation argument like those used in "large matching market" studies (e.g., Roth and Peranson (1999), Kojima and Pathak (2009), Kojima et al. (2013), and Ashlagi et al. (2014)) may prove useful, although this is speculative at this point and beyond the scope of the present paper.

We studied policy goals that we regarded as among the most important ones, but they are far from being exhaustive. Other important policy goals may include a diversity policy requiring certain proportions of different student types in each district (see Nguyen and Vohra (2017) for a related policy at the level of schools), as well as a balanced exchange policy requiring a certain bound on the difference in the numbers of students received from and sent to other districts (see Dur and Ünver (2018) for a related policy at the level

²²Such an example is available from the authors upon request.

of schools). Given that the existing literature has not studied interdistrict school choice, we envision that many policy goals await study within our framework.

While our paper is primarily theoretical and aimed at proposing a general framework to study interdistrict school choice, the main motivation comes from applications to actual programs such as Minnesota's AI program. Given this motivation, it would be interesting to study interdistrict school choice empirically. For instance, evaluating how well the existing programs are doing in terms of balanced exchange, student welfare, and diversity, and how much improvement could be made by a conscious design based on theories such as the ones suggested in the present paper, are important questions left for future work. In addition, implementation of our designs in practice would be interesting. Doing so may, for instance, shed new light on the tradeoff between SPDA and TTC, which has been studied in the intradistrict school choice from a practical perspective (e.g., Abdulkadiroğlu et al. (2006), Abdulkadiroğlu et al. (2017)). We are only beginning to learn about the interdistrict school choice problem, and thus we expect that these and other questions could be answered as more researchers analyze it.

References

- **Abdulkadiroğlu, Atila**, "College admissions with affirmative action," *International Journal of Game Theory*, 2005, 33 (4), 535–549.
- _____ and Tayfun Sönmez, "School choice: A mechanism design approach," *American Economic Review*, June 2003, 93 (3), 729–747.
- _____, Parag A. Pathak, Alvin E. Roth, and Tayfun Sönmez, "The Boston Public School Match," *American Economic Review*, May 2005, 95 (2), 368–371.
 - _____, ____, and _____, "The New York City High School Match," American Economic Review, 2005, 95 (2), 364–367.
- _____, **and** _____, "Strategy-proofness versus Efficiency in Matching with Indifferences: Redesigning the NYC High School Match," *American Economic Review*, 2009, 99, 1954–1978.
 - **_____, Parag Pathak, Alvin E. Roth, and Tayfun Sonmez**, "Changing the Boston School Choice Mechanism," Working Paper 11965, National Bureau of Economic Research January 2006.
- _____, Yeon-Koo Che, Parag A. Pathak, Alvin E. Roth, and Olivier Tercieux, "Minimizing Justified Envy in School Choice: The Design of New Orleans' OneApp," 2017. Working paper.
- **Ahuja, Ravindra K**, *Network flows: theory, algorithms, and applications,* Pearson Education, 2017.

- Aizerman, Mark A. and Andrew V. Malishevski, "General theory of best variants choice: Some aspects," *Automatic Control, IEEE Transactions on*, 1981, 26 (5), 1030–1040.
- Akbarpour, Mohammad and Afshin Nikzad, "Approximate random allocation mechanisms," 2017. Working paper.
- Alkan, Ahmet, "A class of multipartner matching models with a strong lattice structure," *Economic Theory*, June 2002, *19* (4), 737–746.

_____ and David Gale, "Stable schedule matching under revealed preference," *Journal of Economic Theory*, 2003, 112 (2), 289 – 306.

- Ashlagi, Itai, Mark Braverman, and Avinatan Hassidim, "Stability in large matching markets with complementarities," *Operations Research*, 2014, 62 (4), 713–732.
- Aygün, Orhan and Bertan Turhan, "Dynamic Reserves in Matching Markets," Unpublished paper, 2016.
- Aygün, Orhan and Tayfun Sönmez, "Matching with Contracts: Comment," *American Economic Review*, 2013, 103 (5), 2050–2051.
- **Bing, Meir, Daniel Lehmann, and Paul Milgrom**, "Presentation and Structure of Substitutes Valuations," *Proceedings of the 5th ACM Conference on Electronic Commerce*, 2004, pp. 238–239.
- **Biro, P., T. Fleiner, R.W. Irving, and D.F. Manlove**, "The College Admissions Problem with Lower and Common Quotas," *Theoretical Computer Science*, 2010, *411* (34-36), 3136–3153.
- Budish, Eric, Yeon-Koo Che, Fuhito Kojima, and Paul R. Milgrom, "Designing Random Allocation Mechanisms: Theory and Applications," *American Economic Review*, 2013, 103 (2), 585–623.
- **Candogan, Ozan, Markos Epitropou, and Rakesh V Vohra**, "Competitive equilibrium and trading networks: A network flow approach," in "Proceedings of the 2016 ACM Conference on Economics and Computation" ACM 2016, pp. 701–702.
- **Chambers, Christopher P. and M. Bumin Yenmez**, "Choice and Matching," *American Economic Journal: Microeconomics*, August 2017, *9* (3), 126–47.
- _____ **and** _____, "A simple characterization of responsive choice," *Games and Economic Behavior*, 2018, 111, 217 221.
- Che, Yeon-Koo, Jinwoo Kim, and Konrad Mierendorff, "Generalized Reduced-Form Auctions: A Network-Flow Approach," *Econometrica*, 2013, *81* (6), 2487–2520.
- **Clotfelter, Charles T**, "Public school segregation in metropolitan areas.," *Land Economics*, 1999, 75 (4).
- _____, *After Brown: The rise and retreat of school desegregation*, Princeton University Press, 2011.

- **Dur, Umut Mert and M. Utku Ünver**, "Two-sided matching via balanced exchange: Tuition and worker exchanges," 2018. forthcoming, *Journal of Political Economy*.
 - **and Onur Kesten**, "Sequential versus simultaneous assignment systems and two applications," *Economic Theory*, June 2018.
- _____, ____, and M. Utku Ünver, "Flexible Top Trading Cycles and Chains Mechanism: Maintaining Diversity in Erasmus Student Exchange," 2015. Working paper.
 - **_____, Parag Pathak, and Tayfun Sönmez**, "Explicit vs. Statistical Targeting in Affirmative Action: Theory and Evidence from Chicago's Exam Schools," *NBER Working Paper*, 2016, 22109.
- **, Scott D. Kominers, Parag A. Pathak, and Tayfun Sönmez**, "Reserve Design: Unintended Consequences and The Demise of Boston's Walk Zones," 2014. Forthcoming in Journal of Political Economy.
- Echenique, Federico and M. Bumin Yenmez, "How to Control Controlled School Choice," *American Economic Review*, August 2015, *105* (8), 2679–2694.
- Edmonds, Jack and Richard M. Karp, "Theoretical improvements in algorithmic efficiency for network flow problems," *Journal of the ACM (JACM)*, 1972, 19 (2), 248–264.
- Ehlers, Lars, Isa E. Hafalir, M. Bumin Yenmez, and Muhammed A. Yildirim, "School choice with controlled choice constraints: Hard bounds versus soft bounds," *Journal of Economic Theory*, 2014, *153*, 648–683.
- Ekmekci, Mehmet and M. Bumin Yenmez, "Integrating Schools for Centralized Admissions," 2014. Working Paper.
- **Ergin, Haluk and Tayfun Sönmez**, "Games of school choice under the Boston mechanism," *Journal of Public Economics*, 2006, *90* (1-2), 215 237.
- Fragiadakis, Daniel and Peter Troyan, "Improving matching under hard distributional constraints," *Theoretical Economics*, 2017, 12 (2), 863–908.
- _____, Atsushi Iwasaki, Peter Troyan, Suguru Ueda, and Makoto Yokoo, "Strategyproof Matching with Minimum Quotas," ACM Transactions on Economics and Computation, December 2015, 4 (1), 1–40.
- **Fujishige, Satoru and Akihisa Tamura**, "A General Two-sided Matching Market with Discrete Concave Utility Functions," *Discrete Applied Mathematics*, 2006, 154 (6), 950–970.
- and _____, "A Two-sided Discrete-concave Market with Possibly Bounded Side Payments: An Approach by Discrete Convex Analysis," *Mathematics of Operations Research*, 2007, 32 (1), 136–155.
- Gale, David and Lloyd S. Shapley, "College Admissions and the Stability of Marriage," *The American Mathematical Monthly*, January 1962, 69 (1), 9–15.
- Goto, Masahiro, Atsushi Iwasaki, Yujiro Kawasaki, Yosuke Yasuda, and Makotoo

Yokoo, "Improving Fairness and Efficiency in Matching with Distributional Constraints: An Alternative Solution for the Japanese Medical Residency Match," 2014. https://mpra.ub.uni-muenchen.de/53409/1/MPRA_paper_53409.pdf, last accessed on March 9th, 2018.

- _____, Fuhito Kojima, Ryoji Kurata, Akihisa Tamura, and Makoto Yokoo, "Designing Matching Mechanisms under General Distributional Constraints," *American Economic Journal: Microeconomics*, May 2017, 9 (2), 226–62.
- Gul, Faruk, Wolfgang Pesendorfer, and Mu Zhang, "Market Design and Walrasian Equilibrium," *Working Paper*, 2018.
- Hafalir, Isa E., M. Bumin Yenmez, and Muhammed A. Yildirim, "Effective affirmative action in school choice," *Theoretical Economics*, May 2013, *8* (2), 325–363.
- Hassidim, Avinatan, Assaf Romm, and Ran I. Shorrer, "Redesigning the Israeli Psychology Master's Match," *American Economic Review*, May 2017, *107* (5), 205–09.
- Hatfield, John William and Fuhito Kojima, "Matching with contracts: Comment," American Economic Review, 2008, 98 (3), 1189–94.
- _____ **and** _____, "Substitutes and stability for matching with contracts," *Journal of Economic Theory*, 2010, 145 (5), 1704 1723.
- _____ and Paul R. Milgrom, "Matching with Contracts," *American Economic Review*, September 2005, 95 (4), 913–935.
- _____ and Scott Duke Kominers, "Hidden substitutability," 2014. Working Paper.
- Kamada, Yuichiro and Fuhito Kojima, "Efficient Matching under Distributional Constraints: Theory and Applications," *American Economic Review*, 2015, 105 (1), 67–99.
- _____ and _____, "Stability concepts in matching with distributional constraints," *Journal of Economic theory*, 2017, 168, 107–142.
- **and** _____, "Stability and strategy-proofness for matching with constraints: a necessary and sufficient condition," *Theoretical Economics*, 2018, *13* (2), 761–794.
- Kelso, Alexander S. and Vincent P. Crawford, "Job Matching, Coalition Formation, and Gross Substitutes," *Econometrica*, 1982, *50*, 1483–1504.
- Kojima, Fuhito, "School choice: Impossibilities for affirmative action," *Games and Economic Behavior*, 2012, 75 (2), 685 693.
- _____, Akihisa Tamura, and Makoto Yokoo, "Designing matching mechanisms under constraints: An approach from discrete convex analysis," *Journal of Economic Theory*, 2018, 176, 803 833.
- **_____** and Parag A Pathak, "Incentives and stability in large two-sided matching markets," *The American Economic Review*, 2009, pp. 608–627.
- _____, Ning Sun, and Ning Neil Yu, "Job Matching under Constraints," *Working Paper*, 2018.

- _____, **Parag A Pathak, and Alvin E Roth**, "Matching with couples: Stability and incentives in large markets," *The Quarterly Journal of Economics*, 2013, *128* (4), 1585–1632.
- Kominers, Scott Duke and Tayfun Sönmez, "Matching with slot-specific priorities: Theory," *Theoretical Economics*, 2016, 11 (2), 683–710.
- Kurata, Ryoji, Naoto Hamada, Chia-Ling Hsu, Takamasa Suzuki, Suguru Ueda, and Makoto Yokoo, "Pareto efficient strategy-proof school choice mechanism with minimum quotas and initial endowments," in "Proceedings of the 2016 International Conference on Autonomous Agents & Multiagent Systems" International Foundation for Autonomous Agents and Multiagent Systems 2016, pp. 59–67.
- **Manjunath, Vikram and Bertan Turhan**, "Two School Systems, One District: What to Do When a Unified Admissions Process is Impossible," *Games and Economic Behavior*, 2016, *95*, 25 40.
- Milgrom, Paul, "Assignment messages and exchanges," American Economic Journal: Microeconomics, 2009, 1 (2), 95–113.
- _____ and Ilya Segal, "Deferred-acceptance auctions and radio spectrum reallocation.," in "EC" 2014, pp. 185–186.
- **Murota**, Kazuo, *Discrete Convex Analysis*, Society for Industrial and Applied Mathematics, 2003.
- **_____ and Yu Yokoi**, "On the Lattice Structure of Stable Allocations in Two-Sided Discrete-Concave Market," *Mathematics of Operations Research*, 2015, 40, 460–473.
- Nguyen, Thành, Ahmad Peivandi, and Rakesh Vohra, "Assignment problems with complementarities," *Journal of Economic Theory*, 2016, 165, 209–241.
- _____ and Rakesh Vohra, "Stable Matching with Proportionality Constraints," in "Proceedings of the 2017 ACM Conference on Economics and Computation" ACM 2017, pp. 675–676.
- Pycia, Marek and M. Utku Ünver, "Decomposing random mechanisms," *Journal of Mathematical Economics*, 2015, 61, 21–33.
- Rivkin, Steven G., "Residential Segregation and School Integration," Sociology of Education, 1994, 67 (4), 279–292.
- Roth, Alvin E., "Stability and Polarization of Interests in Job Matching," *Econometrica*, January 1984, 52 (1), 47–57.
- **and Elliott Peranson**, "The Redesign of the Matching Market for American Physicians: Some Engineering Aspects of Economic Design," *American Economic Review*, September 1999, *89* (4), 748–780.
- Shapley, Lloyd and Herbert Scarf, "On cores and indivisibility," *Journal of Mathematical Economics*, 1974, 1 (1), 23–37.
- Sönmez, Tayfun, "Bidding for Army Career Specialties: Improving the ROTC Branching

Mechanism," Journal of Political Economy, 2013, 121 (1), 186–219.

- Suzuki, Takamasa, Akihisa Tamura, Naoto Hamada, and Makoto Yokoo, "Efficient allocation mechanism with endowments and distributional constraints," 2017. Working Paper.
- Wells, Amy Stuart, Bianca J Baldridge, Jacquelyn Duran, Courtney Grzesikowski, Richard Lofton, Allison Roda, Miya Warner, and Terrenda White, "Boundary Crossing for Diversity, Equity and Achievement," Technical Report, Harvard Law School 2009.
- Westkamp, Alexander, "An analysis of the German university admissions system," *Economic Theory*, 2013, 53 (3), 561–589.
- Yenmez, M. Bumin, "A college admissions clearinghouse," *Journal of Economic Theory*, 2018, 176, 859 885.

Appendix A. Additional Results

In this section, we provide some additional results. Before we proceed, we introduce two admissions rule properties. An admissions rule Ch satisfies *path independence* if for every $X, Y \subseteq \mathcal{X}$, $Ch(X \cup Y) = Ch(X \cup Ch(Y))$. Path independence states that a set can be divided into not-necessarily disjoint subsets and the admissions rule can be applied to the subsets in any order so that the chosen set of contracts is always the same. An admissions rule Ch satisfies **the irrelevance of rejected contracts** (IRC) if for every $X \subseteq \mathcal{X}$ and $x \notin Ch(X)$, $Ch(X \setminus \{x\}) = Ch(X)$. The irrelevance or rejected contracts states that a rejected contract can be removed from a set without changing the chosen set. Path independence is equivalent to substitutability and IRC (Aizerman and Malishevski, 1981). Furthermore, an admissions rule satisfies substitutability and LAD if, and only if, it satisfies path independence and LAD.²³

A.1. Improving Student Welfare for Centralized Districts. In Section 3.1, we studied when SPDA satisfies individual rationality, which requires that, under interdistrict school choice, every student is matched with a school that is weakly more preferred than her initial school. In this section, we consider an alternative setting where each district uses SPDA to assign its students to schools when there is no interdistrict school choice. In other words, the initial matching is determined by SPDA when there is only intradistrict school choice. More explicitly, each student ranks schools in their home districts (or contracts associated with their home districts) and SPDA is used between a district and students from that district. Note that we assume each student's ranking over contracts associated

²³See Aygün and Sönmez (2013) for a study of IRC and Chambers and Yenmez (2017) for a study of path independence in a matching context.

with the home district is the same as the relative ranking in the original preferences. Importantly, in this setting, the initial matching is not fixed but is determined by student preferences and district admissions rules. In such a setting, we characterize district admissions rules which guarantee that no student is hurt from interdistrict school choice.

The next property of district admissions rules proves to play a crucial role to achieve this policy.

Definition 8. A district admissions rule Ch_d favors own students if for any matching X that is feasible for students,

$$Ch_d(X) \supseteq Ch_d(\{x \in X | d(s(x)) = d\}).$$

When a district admissions rule favors own students, any contract that is chosen from a set of contracts associated with students from a district is also chosen from a superset that includes additional contracts associated with students from the other districts. Roughly, this condition requires that, under interdistrict school choice, a district prioritizes its own students that it used to admit over students from the other districts (even though an out-of-district student can still be admitted when a student from the district is rejected).

The following result shows that this is exactly the condition which guarantees that interdistrict school choice weakly improves the outcome for every student.

Theorem 8. Every student weakly prefers the DA outcome under interdistrict school choice to the DA outcome under intradistrict school choice for all student preferences if, and only if, each district's admissions rule favors own students.

In the proof, we show that in the intradistrict school choice the SPDA outcome can alternatively be produced by an interdistrict school choice model where students rank contracts with all districts and districts have modified admissions rules: For any set of contracts X, each district d chooses the following contracts: $Ch_d(\{x \in X | d(s(x)) = d\})$. Since the original district admissions rules favor own students, the chosen set under the modified admissions rule is a subset of $Ch_d(X)$ when X is feasible for students. Then the conclusion that students receive weakly more preferred outcomes in interdistrict school choice than in intradistrict school choice follows from a comparative statics property of SPDA that we show (Lemma 5).²⁴ To show the "only if" part, when there exists a district admissions rule that fails to favor own students, we construct preferences of students such that interdistrict school choice.

A.2. District-level Type-specific Ceilings. In this section, we show the incompatibility of type-specific ceilings at the district level with the existence of a stable matching.

²⁴We cannot use the comparative statics result of Yenmez (2018) because in our setting $Ch_d(X) \supseteq Ch'_d(X)$ only when X is feasible for students, whereas Yenmez (2018) requires this property for all sets of contracts X.

Definition 9. A district admissions rule Ch_d has a **district-level type-specific ceiling** of q_d^t for type-t students if the number of type-t students admitted cannot exceed this ceiling. More formally, for any matching X that is feasible for students,

$$|\{x \in Ch_d(X) | \tau(s(x)) = t\}| \le q_d^t.$$

Note that, as in the case of school-level type-specific ceilings, district admissions rules do not necessarily satisfy acceptance once district-level type-specific ceilings are imposed. We define a weaker version of the acceptance assumption as follows.

Definition 10. A district admissions rule Ch_d that has district-level type-specific ceilings is *d*weakly acceptant if, for any contract x associated with a type-t student and district d and matching X that is feasible for students, if x is rejected from X, then at $Ch_d(X)$,

- the number of students assigned to school c(x) is equal to $q_{c(x)}$, or
- the number of students assigned to district d is at least k_d , or
- the number of type-t students assigned to district d is at least q_d^t .

This admissions rule property states that a student can be rejected only when one of these three conditions is satisfied.

We establish that in an interdistrict school choice problem in which district admissions rules has district-level type-specific ceilings that also satisfy some other desired properties, there may exist no stable matching.

Theorem 9. There exist districts, schools, students, and their types such that for every admissions rule of a district with district-level type-specific ceilings that satisfies *d*-weak acceptance and IRC, there exist path-independent admissions rules for the other districts and preferences of students such that no stable matching exists.

To show this result, we construct an environment such that a district admissions rule with the desired properties cannot satisfy *weak substitutability*, a necessary condition to guarantee the existence of a stable matching (Hatfield and Kojima, 2008).

Appendix B. Examples of District Admissions Rules

In this section, we first provide a class of district admissions rules that are feasible and acceptant and, furthermore, have completions that satisfy substitutability and LAD. Then, based on this class, we identify admissions rules that also satisfy the properties stated in Theorems 1, 2, 3, and 8.

B.1. An Example of a District Admissions Rule. Consider a district d with schools c_1, \ldots, c_n . Each school c_i has an admissions rule Ch_{c_i} such that, for any set of contracts $X, Ch_{c_i}(X) = Ch_{c_i}(X_{c_i}) \subseteq X_{c_i}$. District d's admissions rule Ch_d is defined as follows. For any set of contracts X,

$$Ch_d(X) = Ch_{c_1}(X) \cup Ch_{c_2}(X \setminus Y_1) \cup \ldots \cup Ch_{c_n}(X \setminus Y_{n-1}),$$

where Y_i for i = 1, ..., n - 1 is the set of all contracts in X associated with students who have contracts in $Ch_{c_1}(X) \cup ... \cup Ch_{c_i}(X \setminus Y_{i-1})$. In words, we order the schools and let schools choose in that order. Furthermore, if a student is chosen by some school, we remove all contracts associated with this student for the remaining schools.

We study when district admissions rule Ch_d satisfies our assumptions.

Claim 1. Suppose that for every school c_i and matching X, $|Ch_{c_i}(X)| \le q_{c_i}$. Then district admissions rule Ch_d is feasible.

Proof. Since every student-school pair uniquely defines a contract, for every matching X, every school c_i , and every student s, there is at most one contract associated with s in $Ch_{c_i}(X)$. In addition, whenever a student's contract with a school c_i is chosen, her contracts with the remaining schools are included in Y_j for every $j \ge i$ by the construction of Ch_d . Hence, for every X, $Ch_d(X)$ is feasible for students. Furthermore, by assumption, $|Ch_{c_i}(X)| \le q_{c_i}$ for each c_i . Therefore, Ch_d is feasible.

Claim 2. Suppose that for every school c_i and matching X, $|Ch_{c_i}(X)| = \min\{q_{c_i}, |X_{c_i}|\}$. Then district admissions rule Ch_d is acceptant.

Proof. Suppose that matching *X* is feasible for students and $x \in X_d \setminus Ch_d(X)$. There exists $i \leq n$ such that $c_i = c(x)$. Since *X* is feasible for students, $x \in X \setminus Y_{i-1}$ where Y_{i-1} is as defined in the construction of Ch_d . Because $x \in X_d \setminus Ch_d(X)$, $x \notin Ch_{c_i}(X \setminus Y_{i-1})$. Then $|Ch_{c_i}(X \setminus Y_{i-1})| = q_{c_i}$ by assumption, which implies that district admissions rule Ch_d is acceptant.

Next we study when district admissions rule Ch_d has a completion that satisfies substitutability and LAD. Consider the following district admissions rule Ch'_d : For any set of contracts X,

$$Ch'_d(X) = Ch_{c_1}(X) \cup \ldots \cup Ch_{c_n}(X).$$

Claim 3. Suppose that for every school c_i , Ch_{c_i} satisfies substitutability and LAD. Then district admissions rule Ch'_d is a completion of Ch_d , and it satisfies substitutability and LAD.

Proof. To show that Ch'_d is a completion of Ch_d , suppose that X is a set of contracts such that $Ch'_d(X)$ is feasible for students. By mathematical induction, we show that $Ch_{c_i}(X) = Ch_{c_i}(X \setminus Y_{i-1})$ for i = 1, ..., n, where Y_i is defined as above for i > 1 and $Y_0 = \emptyset$. The claim trivially holds for i = 1. Suppose that it also holds for 1, ..., i - 1. We show the claim for i. Since $Ch'_d(X)$ is feasible for students, $Ch_{c_i}(X)$ and $Ch_{c_1}(X) \cup ... \cup Ch_{c_{i-1}}(X)$ do not have any contracts associated with the same student. Therefore, $Ch_{c_i}(X) \cap Y_{i-1} = \emptyset$. Since Ch_{c_i} satisfies IRC, $Ch_{c_i}(X) = Ch_{c_i}(X \setminus Y_{i-1})$. As a result, $Ch_d(X) = Ch'_d(X)$, which completes the proof that Ch'_d is a completion of Ch_d .

37

Since all school admissions rules satisfy substitutability and LAD, so does Ch'_d .

All of the assumptions on school admissions rules stated in Claims 1, 2, and 3 are satisfied when school admissions rules are *responsive*: each school has a ranking of contracts associated with itself and, from any given set of contracts, each school chooses contracts with the highest rank until the capacity of the school is full or there are no more contracts left. Responsive admissions rules satisfy substitutability and LAD. Furthermore, for every school c_i , $|Ch_{c_i}(X)| = \min\{q_{c_i}, |X_{c_i}|\}$.²⁵ By the claims stated above, when school admissions rules are responsive, district admissions rule Ch_d is feasible and acceptant, and it has a completion that satisfies substitutability and LAD.

Based on these results, we provide examples of district admissions rules that further satisfy the additional assumptions considered in different parts of our paper.

B.2. District Admissions Rules Satisfying the Assumptions in Theorem 1. We use the district admissions rule construction above and we further specify each school's admissions rule. Each school has a responsive admissions rule. If a student is initially matched with a school, then her contract with this school is ranked higher than contracts of students who are not initially matched with the school. As before, district admissions rule Ch_d is feasible and acceptant, and it has a completion that satisfies substitutability and LAD.

Claim 4. District admissions rule Ch_d respects the initial matching.

Proof. Let *c* be the initial school of student *s* and x = (s, d, c). By construction, for any matching *X* that is feasible for students, $x \in X$ implies $x \in Ch_d(X)$ because *c* chooses *x* from any set of contracts and *s* does not have any other contract in *X*. Therefore, Ch_d respects the initial matching.

B.3. District Admissions Rules Satisfying the Assumptions in Theorem 2. We modify the district admissions rule construction in Appendix B.1. Each school has a ranking of contracts associated with itself. When it is the turn of a school, it accepts contracts that have the highest rank until the capacity of the school is full, or the number of contracts chosen by the district is k_d , or there are no more contracts left. The remaining contracts of a chosen student are removed.

District admissions rule Ch_d is feasible because no school admits more students than its capacity and no student is admitted to more than one school.

Claim 5. *District admissions rule* Ch_d *is acceptant.*

²⁵See Chambers and Yenmez (2018) for a characterization of responsive admissions rules using substitutability.

Proof. To show acceptance, suppose that matching X is feasible for students and $x \in X_d \setminus Ch_d(X)$. There exists $i \leq n$ such that $c_i = c(x)$. Since X is feasible for students, $x \in X \setminus Y_{i-1}$ where Y_{i-1} is the set of all contracts in X associated with students who are chosen by schools c_1, \ldots, c_{i-1} . Because $x \in X_d \setminus Ch_d(X)$, x is not chosen by c_i . Then, by construction, either c_i fills its capacity or the district admits k_d students, which implies that Ch_d is acceptant.

Claim 6. District admissions rule Ch_d has a completion that satisfies substitutability and LAD.

Proof. First, we construct a completion of Ch_d . Define the following district admissions rule: given a set of contracts X, when it is the turn of a school, it chooses from all the contracts in X. Each school chooses contracts using the same priority order until the school capacity is full, or the district has k_d contracts, or there are no more contracts left. Denote this admissions rule by Ch'_d . Suppose that $Ch'_d(X)$ is feasible for students. Then, by construction, $Ch'_d(X) = Ch_d(X)$. Therefore, Ch'_d is a completion of Ch_d .

Next, we show that Ch'_d satisfies LAD. Suppose that $Y \supseteq X$. Every school c_i chooses weakly more contracts from Y than X unless the number of contracts chosen from Y by the district reaches k_d . Since the number of chosen contracts from X cannot exceed k_d by construction, Ch'_d satisfies LAD.

Finally, we show that Ch'_d satisfies substitutability. Suppose that $x \in X \subseteq Y$ and $x \in Ch'_d(Y)$. Therefore, the number of contracts chosen from Y by schools preceding c(x) is strictly less than k_d . This implies that the number of contracts chosen from X by schools preceding c(x) is weakly less than this number as weakly more contracts are chosen by schools preceding school c(x) in Y than X. As a result, for school c(x), weakly more contracts can be chosen from X than Y.

The ranking of contract x among Y in the ranking of school c(x) is high enough that it is chosen from set Y. Therefore, the ranking of contract x among X in the ranking of school c(x) must be high enough to be chosen from set X because weakly more contracts are chosen from X than Y for school c(x).

Furthermore, by construction, district admissions rule Ch_d never chooses more than k_d students. Therefore, it is also rationed.

B.4. District Admissions Rules Satisfying the Assumptions in Theorem 3. District admissions rules can accommodate unmatched students by *reserving* seats for different types of students:

Definition 11. Let c be a school in district d. A district admissions rule Ch_d has a reserve of r_c^t for type-t students at school c if, for any feasible matching X that does not have any contract associated with type-t student s, if $|\{x \in X_c | \tau(s(x)) = t\}| < r_c^t$, then x = (s, d, c) satisfies $x \in Ch_d(X \cup \{x\})$.

A reserve for a student type at a school *c* guarantees space for this type at school *c*. Therefore, when a student is unmatched at a feasible matching and the reserve for her type is not yet filled at a school, the district will accept this student at that school if she applies to it.

Claim 7. Suppose that districts have admissions rules with reserves such that $\sum_{c} r_{c}^{t} = k^{t}$ for every type t. Then district admissions rules accommodate unmatched students.

Proof. Suppose that student *s* is unmatched at a feasible matching *X*. Let *t* be the type of student *s*. Then there exists a school *c* such that the number of type-*t* students in *c* at *X* is strictly less than r_c^t because $\sum_c r_c^t = k^t$. By definition of reserves, x = (s, c) satisfies $x \in Ch_{d(c)}(X \cup \{x\})$.

A district can have type-specific reserves at its schools in different ways. In the rest of this subsection, we use school admissions rules with reserves introduced by Hafalir et al. (2013) to construct a fairly general example in which a district has schools with type-specific reserves. Let r_c^t be the number of seats reserved by school c for type-t students. Suppose that the type-specific ceilings for schools are given and that they satisfy the assumptions in Section 3.3. Assume that, for every district d, $\sum_{c:d(c)=d} \sum_t r_c^t = k_d$, $\sum_c r_c^t = k^t$ and, for every type t and school c, $r_c^t \leq q_c^t$. Furthermore, assume that $\sum_t r_c^t \leq q_c$ for every school c.

Consider the following district admissions rule for district d. Schools are ordered as c_1, c_2, \ldots, c_n . Each school has a ranking over contracts associated with it and a linear order over student types. First, all schools choose contracts for their reserve seats according to the order c_1, c_2, \ldots, c_n . When it is the turn of school c_i , all contracts associated with students whose contracts were previously chosen are removed. School c_i chooses contracts for its reserved seats so that, for every type, either reserved seats are filled or there are no more contracts associated with students of that type remaining. Then all schools choose contracts of previously chosen students are removed. School c_i chooses from the remaining contracts in order. When a contract of a type-t student is considered, this contract is chosen unless the school's capacity is filled or its type-t ceiling is filled or the district has k_d contracts. Denote this district admissions rule by Ch_d .

District admissions rule Ch_d is feasible because a student cannot have more than one contract and a school cannot have more contracts than its capacity at any chosen set of contracts. It is also weakly acceptant and rationed by construction. Furthermore, for every type t and school c, the district cannot admit more than q_c^t type-t students at c, so it has a school-level type-specific ceiling of q_c^t for type-t students and school c.

Claim 8. District admissions rule Ch_d has a completion that satisfies substitutability and LAD.

Proof. For any set of contracts X, school c, and type t, let X_c^t denote the set of all contracts in X that are associated with school c and type-t students.

Consider the construction of Ch_d above, but modify it by not removing contracts of students who are chosen previously. Denote this district admissions rule by Ch'_d . To show that Ch'_d is a completion of Ch_d , consider a set of contracts X and suppose that $Ch'_d(X)$ is feasible for students. Since the only difference in the constructions of Ch_d and Ch'_d is the removal of contracts of previously chosen students, it must be that $Ch'_d(X) = Ch_d(X)$. Therefore, Ch'_d is a completion of Ch_d .

To prove substitutability of Ch'_d , suppose, for contradiction, that there exist sets of contracts X and Y with $X \subseteq Y$ and a contract $x \in X$ such that $x \in Ch'_d(Y) \setminus Ch'_d(X)$. Let sand c be such that x = (s, c) and $t = \tau(s)$. First, note that $|X_c^t| > r_c^t$ because $x \notin Ch'_d(X)$. Since $Y \supseteq X$, $|Y_c^t| \ge |X_c^t| > r_c^t$ is implied. Therefore, it is after all schools in d have chosen contracts based on their reserves in the algorithm describing Ch'_d that contract x is chosen by Ch'_d given Y. Let n(X) and n(Y) be the numbers of contracts that have been chosen by all schools before the step (call it step κ_c) at which school c chooses students beyond its reserve under X and Y, respectively. Because $x \in Ch'_d(Y)$, it follows that $n(Y) < k_d$. Therefore, for each school c', the number of contracts chosen by c' before step κ_c under Yis weakly larger than those under X, which we prove as follows:

- Suppose that school c' is processed after school c in the algorithm deciding Ch_d . Then, by step κ_c , c' is matched with students of each type t' only up to its type-t' reserve. More formally, the numbers of type-t' students matched to c' are equal to $\min\{r_{c'}^{t'}, |X_{c'}^{t'}|\}$ and $\min\{r_{c'}^{t'}, |Y_{c'}^{t'}|\}$ under X and Y, respectively. Obviously, the latter expression is no smaller than the former expression.
- Suppose that school c' is processed before school c in the algorithm deciding Ch_d . Recall that $n(Y) < k_d$. Therefore, for school c', it is either (i) as many as $q_{c'}$ students are matched to c' under $Ch_d(Y)$, or (ii) for each type t', the number of type-t' students matched to c' in Y is $\min\{q_{c'}^{t'}, |Y_{c'}^{t'}|\}$. In case (i), the desired conclusion follows trivially because, given any set of contracts, the number of students matched to c' cannot exceed $q_{c'}$. For case (ii), under X, the number of type-t' students matched to c' cannot exceed $\min\{q_{c'}^{t'}, |X_{c'}^{t'}|\} \le \min\{q_{c'}^{t'}, |Y_{c'}^{t'}|\}$. Summing up across all types, we obtain the desired conclusion.

Thus $n(X) \leq n(Y)$, so $k_d - n(X) \geq k_d - n(Y)$. Now, in step κ_c , school c will choose all the applications until either the total number of contracts chosen reaches k_d , or the total number of contracts chosen at c reaches q_c , or the number of contracts chosen at c that are associated with type t students reaches q_c^t . Given the previous fact that $k_d - n(X) \geq k_d - n(Y)$, the fact that $Y \supseteq X$, and the fact that x is chosen by c in step κ_c under Y, it has to be the case that x is also chosen by c under X in step κ_c or before. We prove this as follows:

- If $|X_c^t| \le r_{c'}^t$ then x is chosen in the reserve stage by construction.
- Let $|X_c^t| > r_c^t$. First note that at step κ_c under X and Y, for each type t, there are fewer contracts associated with school c and type-t students that remain to be processed under X than under Y ($X \subseteq Y$, and there is no contract in $X = X \cap Y$ that is processed in the reserve stage under Y but not under X), so the subset of X that should be processed in κ_c is a subset of the corresponding subset of Y. Moreover, the remaining number of contracts to be chosen before reaching the ceiling at c for each type t in step κ_c is weakly larger at X than at Y by the definition of the reserve stage. Finally, as argued above, the total number of students in the district who can still be chosen at κ_c is weakly larger under X than at Y, so whenever x is chosen under Y in this stage, x is chosen under X in this stage or the reserve stage.

This is a contradiction to the assumption that $x \notin Ch'_d(X)$.

To show that Ch'_d satisfies LAD, suppose, for contradiction, that there exist two sets of contracts X, Y with $X \subseteq Y$ and $|Ch'_d(Y)| < |Ch'_d(X)|$. Then, because $Ch'_d(Y) = \bigcup_{c:d(c)=d} (Ch'_d(X) \cap Y_c)$ and $Ch'_d(X) = \bigcup_{c:d(c)=d} (Ch'_d(X) \cap X_c)$, there exists a school c with d(c) = d such that

(1)
$$|Ch'_d(Y) \cap Y_c| < |Ch'_d(X) \cap X_c|$$

Fix such *c* arbitrarily. Next, note that

$$|Ch'_d(Y)| < |Ch'_d(X)| \le k_d,$$
$$|Ch'_d(Y) \cap Y_c| < |Ch'_d(X) \cap X_c| \le q_c,$$

where the first line follows because Ch'_d is rationed by construction, and the second line also holds by construction of Ch'_d . Therefore,

(2)
$$\begin{aligned} |Ch'_{d}(Y) \cap Y^{t}_{c}| &= \min\{|Y^{t}_{c}|, q^{t}_{c}\}\\ &\geq \min\{|X^{t}_{c}|, q^{t}_{c}\}\\ &= |Ch'_{d}(X) \cap X^{t}_{c}|, \end{aligned}$$

for each type $t \in \mathcal{T}$. Because $Ch'_d(Y) \cap Y_c = \bigcup_{t \in \mathcal{T}} (Ch'_d(Y) \cap Y_c^t)$ and $Ch'_d(X) \cap X_c = \bigcup_{t \in \mathcal{T}} (Ch'_d(X) \cap X_c^t)$, inequality (2) and the fact $Y_c^t \cap Y_c^{t'} = X_c^t \cap X_c^{t'} = \emptyset$ for any pair of types t, t' with $t \neq t'$ imply

$$|Ch'_d(Y) \cap Y_c| \ge |Ch'_d(X) \cap X_c|,$$

which contradicts inequality (1).

41

B.5. District Admissions Rules Satisfying the Assumptions in Theorem 8. Consider the district admissions rule construction in Appendix B.1. In this example, let each school use a priority ranking in such a way that all contracts of students from district *d* are ranked higher than the other contracts.

Claim 9. District admissions rule Ch_d favors own students.

Proof. Suppose that X is feasible for students. When it is the turn of school c_i , it considers X_{c_i} . Therefore, $Ch_d(X) = Ch_{c_1}(X_{c_1}) \cup \ldots \cup Ch_{c_k}(X_{c_k})$. Furthermore, $Ch_{c_i}(X_{c_i}) \supseteq Ch_{c_i}(\{x \in X_{c_i} | d(s(x)) = d\})$ by construction. Taking the union of all subset inclusions yields $Ch_d(X) \supseteq Ch_d(\{x \in X_d | d(s(x)) = d\})$. Therefore, Ch_d favors own students. \Box

Appendix C. An Example for Diversity

In this section, we provide an example in which the conditions on the admissions rules stated in Theorem 3 are satisfied and, therefore, SPDA satisfies the diversity policy.

Consider a problem with two school districts, d_1 and d_2 . District d_1 has school c_1 with capacity three and school c_2 with capacity two. District d_2 has school c_3 with capacity two and school c_4 with capacity one. There are seven students: students s_1 , s_2 , s_3 , and s_4 are from district d_1 , whereas students s_5 , s_6 , and s_7 are from district d_2 . Students s_1 , s_5 , s_6 , and s_7 have type t_1 and s_2 , s_3 , and s_4 have type t_2 . To construct district admissions rules that satisfy the properties stated in Theorem 3, let us first specify type-specific ceilings and calculate implied floors and implied ceilings. Suppose that

$$\begin{aligned} q_{c_1}^{t_1} &= 1, \; q_{c_1}^{t_2} = 2, \; q_{c_2}^{t_1} = 1, \; q_{c_2}^{t_2} = 1, \\ q_{c_3}^{t_1} &= 2, \; q_{c_3}^{t_2} = 1, \; q_{c_4}^{t_1} = 1, \; q_{c_4}^{t_2} = 1. \end{aligned}$$

These yield the following implied floors,²⁶

$$\hat{p}_{d_1}^{t_1} = 1, \ \hat{p}_{d_1}^{t_2} = 2, \ \hat{p}_{d_2}^{t_1} = 2, \ \hat{p}_{d_2}^{t_2} = 0,$$

and implied ceilings

$$\hat{q}_{d_1}^{t_1} = 2, \ \hat{q}_{d_1}^{t_2} = 3, \ \hat{q}_{d_2}^{t_1} = 3, \ \hat{q}_{d_2}^{t_2} = 1.$$

²⁶ To see this, note that there cannot be zero type- t_1 students in d_1 (otherwise not all type- t_1 students can be matched since there are only three spaces available for type- t_1 students in d_2). If there is one type- t_1 student in d_1 , there has to be three type- t_1 students in d_2 , which implies there cannot be any type- t_2 students in d_2 , and this implies there will be three type- t_2 students in d_1 . If there are two type- t_1 students in d_1 , there have to be two type- t_2 students in d_2 , which implies there is one type- t_2 student in d_2 , and this implies there will be three type- t_2 students and maximum numbers, we obtain the implied reserves and implied ceilings accordingly. These bounds are achievable because it is feasible to have (i) one type- t_1 student in d_1 , three type- t_1 students in d_2 , zero type- t_2 students in d_2 , and three type- t_2 students in d_1 , and (ii) two type- t_1 students in d_1 , two type- t_2 students in d_2 , one type- t_2 student in d_2 , and two type- t_2 students in d_1 .

For any type *t* and two districts *d* and *d'*, denote $\hat{q}_{d}^{t}/k_{d} - \hat{p}_{d'}^{t}/k_{d'}$ by $\Delta_{d,d'}^{t}$. Using the implied floors and ceilings above, we get:

$$\begin{split} \Delta_{d_1,d_2}^{t_1} &= 2/4 - 2/3 = -1/6, \\ \Delta_{d_2,d_1}^{t_1} &= 3/3 - 1/4 = 3/4, \\ \Delta_{d_1,d_2}^{t_2} &= 3/4 - 0/3 = 3/4, \text{ and} \\ \Delta_{d_2,d_1}^{t_2} &= 1/3 - 2/4 = -1/6. \end{split}$$

Hence, these type-specific ceilings satisfy the condition stated in Theorem 3 that $\Delta_{d,d'}^t \leq \alpha$ for $\alpha = 0.75$.

We construct district admissions rules that have type-specific ceilings, accommodate unmatched students, and are rationed and weakly acceptant. As in Appendix B.4, we consider type-specific reserves (as detailed below, we first fill in the reserves while applying the district admissions rule that uses type-specific reserves). Let us consider the reserves for schools as follows:

$$r_{c_4}^{t_2} = 0$$
, and $r_c^t = 1$ for all other c, t

Consider the following district admissions rule. For each district, schools and student types are ordered and each school a linear order over students. First, schools choose contracts for their reserved seats following the master priority list until the reserves are filled or all the applicants of the relevant type are processed. Then, following the given order over schools and student types, schools choose from the remaining contracts following the linear order over students in order to fill the rest of their seats until the school capacity is filled, or the district has k_d contracts, or district type-specific ceilings are filled, or there are no more remaining contracts.²⁷

To give a more concrete example, suppose that the linear order over students for each school is as follows: $s_1 \succ s_2 \succ s_3 \succ s_4 \succ s_5 \succ s_6 \succ s_7$ and schools and types are ordered from the lowest index to the highest. Then, for example, we have the following:

$$Ch_{d_1}(\{(s_1, c_1), (s_2, c_1), (s_3, c_1), (s_4, c_1), (s_5, c_2), (s_6, c_2)\}) = \{(s_1, c_1), (s_2, c_1), (s_3, c_1), (s_5, c_2)\}.$$

Let us elaborate on how we determine the chosen set of contracts in the above case. School c_1 considers contracts with students s_1 , s_2 , s_3 , and s_4 . Among these students, c_1 accepts s_1 for its reserve for type t_1 , and s_2 for its reserve for type t_2 . Moreover, school c_2 considers contracts with students s_5 and s_6 . Among these students students, c_2 accepts s_5 for its reserve for type t_2 . For the remainder of seats, s_3 is accepted by c_1 since (i) c_1 's type t_2 ceiling is not full, (ii) c_1 's capacity is not full, and (iii) district d_1 has only three accepted

²⁷In Appendix B.4, we provide a class of admissions rules that include the one we consider here. These admissions rules satisfy all of the assumptions that we make in this section.

contracts at this point. Next, s_4 and s_5 are rejected since d_1 has accepted four contracts at this point. This results in the chosen set of contracts presented above.

To illustrate the SPDA outcomes, consider student preferences given by the following table.

$\underline{P_{s_1}}$	$\underline{P_{s_2}}$	P_{s_3}	$\underline{P_{s_4}}$	P_{s_5}	$\underline{P_{s_6}}$	P_{s_7}
c_2	C_3	c_4	c_1	c_1	c_1	c_2
:	c_1	c_2	c_3	c_2	c_4	c_3
	÷	÷	c_2	c_3	c_3	÷
			c_4	c_4	c_2	

SPDA results in the following outcome:

$$\{(s_1, c_2), (s_2, c_3), (s_3, c_2), (s_4, c_1), (s_5, c_1), (s_6, c_4), (s_7, c_3)\}.$$

District d_1 gets two students of both types and district d_2 gets two type- t_1 students and one type- t_2 student. As a result, the ratio difference for type- t_1 students between these districts is roughly 0.17, and the ratio difference for type- t_2 students is roughly 0.17. This example illustrates that the actual ratio differences can be significantly lower than the one given by Theorem 3 (0.17 versus 0.75).

Appendix D. Omitted Proofs

In this section, we provide the omitted proofs.

Proof of Theorem 1. First, to show the "if" part, suppose that all district admissions rules respect the initial matching. In SPDA, each student *s* goes down in her preference order, and either SPDA ends before student *s* reaches her initial school (which is a preferred outcome over the initial school), or student *s* reaches her initial school. In the latter case, she is matched with her initial school because the district's admissions rule respects the initial matching and the district always considers a set of contracts that is feasible for students at any step of SPDA. From this step on, the district accepts this contract, so student *s* is matched with her initial school. Therefore, SPDA satisfies individual rationality.

To prove the "only if" part, suppose that there exists a district d with an admissions rule that fails to respect the initial matching. Hence, there exists a matching X, which is feasible for students, that includes x = (s, d, c) where school c is the initial school of student s and $x \notin Ch_d(X)$. Now, consider student preferences such that every student associated with a contract in X_d prefers that contract the most and all other students prefer a contract associated with a different district the most. Then, at the first step of SPDA, district d considers matching X_d and tentatively accepts $Ch_d(X_d)$. Since $x \notin Ch_d(X_d)$, contract x is rejected at the first step. Therefore, student s is matched with a strictly less preferred school than her initial school, which implies that SPDA does not satisfy individual rationality.

Proof of Theorem 2. We first prove that if each district admissions rule is rationed, then SPDA satisfies the balanced-exchange policy. Let X be the matching produced by SPDA for a given preference profile.

We begin by showing that each student must be matched with a school in X. Suppose, for contradiction, that student s is unmatched. Since X is a stable matching, every contract x = (s, d, c) associated with the student is rejected by the corresponding district, i.e., $x \notin Ch_d(X \cup \{x\})$. Otherwise, student s and district d would like to match with each other using contract x, contradicting the stability of matching X. Since $X \cup \{x\}$ is feasible for students, acceptance implies that, for each district d, either every school in the district is full or that the district has at least k_d students at matching X. Both of them imply that the district has at least k_d . But this is a contradiction to the assumption that student s is unmatched since the existence of an unmatched student implies that there is at least one district d such that the number of students in X_d is less than k_d . Therefore, all students are matched in X.

Because *X* is the outcome of SPDA, it is feasible for students. Therefore, because district admissions rules are rationed, the number of students in district *d* cannot be strictly more than k_d for any district *d*. Furthermore, since every student is matched, the number of students in district *d* must be exactly k_d (because, otherwise, at least one student would have been unmatched.) As a result, SPDA satisfies the balanced-exchange policy.

Next, we prove that if at least one district's admissions rule fails to be rationed, then there exists a student preference profile under which SPDA does not satisfy the balancedexchange policy. Suppose that there exist a district d and a matching X, which is feasible for students, such that $|Ch_d(X)| > k_d$. Consider a feasible matching X' such that (i) all students are matched, (ii) $X'_d = Ch_d(X)$, and (iii) for every district $d' \neq d$, $|X'_{d'}| \leq k_{d'}$. The existence of such X' is guaranteed since every district has enough capacity to serve its students (i.e., for every district d', $\sum_{c:d(c)=d'} q_c \geq k_{d'}$), and $|Ch_d(X)| > k_d$. Now, consider any student preferences, where every student likes her contract in X' the most.

We show that SPDA stops in the first step. For district $d' \neq d$, $X'_{d'}$ is feasible and the number of students matched to d' at $X'_{d'}$ is weakly less than $k_{d'}$. Since $Ch_{d'}$ is acceptant, $Ch_{d'}(X'_{d'}) = X'_{d'}$. For district d, we need to show that $Ch_d(X'_d) = X'_d$, which is equivalent to $Ch_d(Ch_d(X)) = Ch_d(X)$. Let Ch'_d be a completion of Ch_d that satisfies path independence. Because X and $Ch_d(X)$ are feasible for students, $Ch'_d(X) = Ch_d(X)$ and $Ch'_d(Ch'_d(X)) = Ch_d(Ch_d(X))$. Furthermore, since Ch'_d is path independent, $Ch'_d(Ch'_d(X)) = Ch'_d(X)$, which implies $Ch_d(Ch_d(X)) = Ch_d(X)$. As a result, $Ch_d(X'_d) = X'_d$. Therefore, SPDA stops at the first step since no contract is rejected.

Since SPDA stops at the first step, the outcome is matching X'. But X' fails the balanced-exchange policy because $|X'_d| = |Ch_d(X)| > k_d$.

Proof of Theorem 3. To prove this result, we provide the following lemmas.

Lemma 2. *If district admissions rules accommodate unmatched students, every student is matched to a school in SPDA.*

Proof of Lemma 2. Let *X* be the outcome of SPDA for some preference profile. Suppose, for contradiction, that student *s* is unmatched. Since *X* is a stable matching and student *s* prefers any contract x = (s, d, c) to being unmatched, $x \notin Ch_d(X \cup \{x\})$. But this is a contradiction to the assumption that district admissions rules accommodate unmatched students.

Lemma 3. For each type t, district d, and legitimate matching X, we have $\hat{q}_d^t \ge \xi_d^t(X) \ge \hat{p}_d^t$. Moreover, for each type t and district d, there exists a legitimate matching X such that $\xi_d^t(X) = \hat{p}_d^t$ and there exists a legitimate matching X such that $\xi_d^t(X) = \hat{q}_d^t$.

Proof of Lemma 3. Observe that for every legitimate matching *X*, the induced distribution satisfies the constraints of the linear program. Therefore, the first part follows from the definition of the implied floors and ceilings. For the second part, note that there exists a solution to the linear program such that the ceiling and the floor are attained. Furthermore, every solution $y = (y_c^t)_{c \in C, t \in T}$ of the linear program can be supported by a legitimate matching *X* such that $y_c^t = \xi_c^t(X)$ for every *c* and *t*.

Lemma 4. For each $t \in \mathcal{T}$ and $d, d' \in \mathcal{D}$ with $d \neq d'$, there exists a legitimate matching X such that $\xi_d^t(X) = \hat{q}_d^t$ and $\xi_{d'}^t(X) = \hat{p}_{d'}^t$.

Proof of Lemma 4. Let \hat{X} be a legitimate matching such that $\xi_d^t(\hat{X}) = \hat{q}_d^t$ and \mathcal{M}_0 be the set of all legitimate matchings. Let

$$\mathcal{M}_1 \equiv \{ X \in \mathcal{M}_0 | \xi_{d'}^t(X) = \hat{p}_{d'}^t \}.$$

Note that M_1 is nonempty due to Lemma 3. Next, let

$$\mathcal{M}_2 \equiv \{ X \in \mathcal{M}_1 | \sum_{\tilde{t},\tilde{c}} | \xi_{\tilde{c}}^{\tilde{t}}(X) - \xi_{\tilde{c}}^{\tilde{t}}(\hat{X}) | \leq \sum_{\tilde{t},\tilde{c}} | \xi_{\tilde{c}}^{\tilde{t}}(X') - \xi_{\tilde{c}}^{\tilde{t}}(\hat{X}) | \text{ for every } X' \in \mathcal{M}_1 \}.$$

Note that \mathcal{M}_2 is nonempty because \mathcal{M}_1 is a finite set. We will show that for any $X \in \mathcal{M}_2$, $\xi_d^t(X) = \xi_d^t(\hat{X}) = \hat{q}_d^t$.

47

To prove the above claim, assume for contradiction that there exists $X \in \mathcal{M}_2$ such that $\xi_d^t(X) \neq \xi_d^t(\hat{X})$. By Lemma 3, $\xi_d^t(X) \neq \xi_d^t(\hat{X})$ implies that $\xi_d^t(X) < \xi_d^t(\hat{X})$. Then there exists c with d(c) = d such that $\xi_c^t(X) < \xi_c^t(\hat{X})$. Consider the following procedure.

Step 0: Initialize by setting $(t_1, c_1) := (t, c)$. Note that $\xi_{c_1}^{t_1}(X) < \xi_{c_1}^{t_1}(\hat{X})$ by definition of *c*.

Step $i \ge 1$: We begin with (t_i, c_i) . Note that (by assumption for i = 1, and as shown later for $i \ge 2$), $\xi_{c_i}^{t_i}(X) < \xi_{c_i}^{t_i}(\hat{X})$. Denote $d_i = d(c_i)$. Now,

- (1) Suppose that there exists i' < i such that either (i) $c_{i'}^* = c_i$ or (ii) $\xi_{c_i}^{t_i}(X) < q_{c_i}$ and $d(c_{i'}^*) = d(c_i)$. If such an index i' exists, then set $(t_{i+1}, c_i^*) := (t_{i'+1}, c_{i'}^*)$.
- (2) Suppose not. Then, if there exists $t' \in \mathcal{T}$ such that $\xi_{c_i}^{t'}(X) > \xi_{c_i}^{t'}(\hat{X})$, then set $t_{i+1} := t'$ and $c_i^* := c_i$.
- (3) If not, then note that ∑_{t∈T} ξ^t_{ci}(X) < q_{ci}.²⁸ Also note that there exists a type-school pair (t', c') with c' ≠ c_i such that ξ^{t'}_{ci}(X) > ξ^{t'}_{ci}(X̂) and d(c') = d_i because ∑_{c̄:d(c̄)=d_i,t̄∈T} ξ^t_{c̄}(X) = ∑_{c̄:d(c̄)=d_i,t̄∈T} ξ^t_{c̄}(X̂) = k_{di}.
 (a) If t' = t_i, then let X̃ be a matching such that

a) If $t = t_i$, then let A be a matching such that

$$\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \begin{cases} \xi_{c_i}^{t_i}(X) + 1 & \text{ for } (\tilde{t}, \tilde{c}) = (t_i, c_i), \\ \xi_{c'}^{t'}(X) - 1 & \text{ for } (\tilde{t}, \tilde{c}) = (t_i, c'), \\ \xi_{\tilde{c}}^{\tilde{t}}(X) & \text{ otherwise.} \end{cases}$$

Note that $\tilde{X} \in \mathcal{M}_1$.²⁹ Also, by construction, $\sum_{\tilde{t},\tilde{c}} |\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) - \xi_{\tilde{c}}^{\tilde{t}}(\hat{X})| = \sum_{\tilde{t},\tilde{c}} |\xi_{\tilde{c}}^{\tilde{t}}(X) - \xi_{\tilde{c}}^{\tilde{t}}(\hat{X})| - 2 < \sum_{\tilde{t},\tilde{c}} |\xi_{\tilde{c}}^{\tilde{t}}(X) - \xi_{\tilde{c}}^{\tilde{t}}(\hat{X})|$, which contradicts the assumption that $X \in \mathcal{M}_2$.

- (b) Therefore, suppose that $t' \neq t_i$ and let $t_{i+1} := t'$ and $c_i^* := c'$.
- (4) The pair (t_{i+1}, c_i^*) created above satisfies $\xi_{c_i^*}^{t_{i+1}}(X) > \xi_{c_i^*}^{t_{i+1}}(\hat{X})$, so there exists $c' \in \mathcal{C}$ such that $\xi_{c'}^{t_{i+1}}(X) < \xi_{c'}^{t_{i+1}}(\hat{X})$. Set $c_{i+1} = c'$. Note that $\xi_{c_{i+1}}^{t_{i+1}}(X) < \xi_{c_{i+1}}^{t_{i+1}}(\hat{X})$.

We follow the procedure above to define $(t_1, c_1), (t_2, c_1^*), (t_2, c_2), (t_3, c_2^*), (t_3, c_3)$, and so forth. Because the set \mathcal{T} is finite, we have *i* and j > i with $t_i = t_j$. Consider the smallest *j* with this property (note that given such *j*, *i* is uniquely identified). Now, let \tilde{X} be a

²⁸A proof of this fact is as follows. By an earlier argument, $\xi_{c_i}^{t_i}(X) < \xi_{c_i}^{t_i}(\hat{X})$. Moreover, by assumption $\xi_{c_i}^{\tilde{t}}(X) \leq \xi_{c_i}^{\tilde{t}}(\hat{X})$ for every $\tilde{t} \in \mathcal{T}$. Therefore, $\sum_{\tilde{t} \in \mathcal{T}} \xi_{c_i}^{\tilde{t}}(X) < \sum_{\tilde{t} \in \mathcal{T}} \xi_{c_i}^{\tilde{t}}(\hat{X}) \leq q_{c_i}$. ²⁹A proof of this fact is as follows. Because $\sum_{\tilde{t} \in \mathcal{T}} \xi_{c_i}^{\tilde{t}}(X) < q_{c_i}, \sum_{\tilde{t} \in \mathcal{T}} \xi_{c_i}^{\tilde{t}}(\hat{X}) = \sum_{\tilde{t} \in \mathcal{T}} \xi_{c_i}^{\tilde{t}}(X) + 1 \leq q_{c_i}$. For

²⁹A proof of this fact is as follows. Because $\sum_{\tilde{t}\in\mathcal{T}}\xi_{c_i}^t(X) < q_{c_i}, \sum_{\tilde{t}\in\mathcal{T}}\xi_{c_i}^t(X) = \sum_{\tilde{t}\in\mathcal{T}}\xi_{c_i}^t(X) + 1 \le q_{c_i}$. For every $\tilde{c} \neq c_i, \sum_{\tilde{t}\in\mathcal{T}}\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) \le \sum_{\tilde{t}\in\mathcal{T}}\xi_{\tilde{c}}^{\tilde{t}}(X) \le q_{\tilde{c}}$. Thus, all school capacities are satisfied. For all $\tilde{c}, \tilde{t}, \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) \le \max\{\xi_{\tilde{c}}^{\tilde{t}}(X), \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X})\} \le q_{\tilde{c}}^{\tilde{t}}$ by construction, so all type-specific ceilings are satisfied. And $\sum_{\tilde{t}\in\mathcal{T},\tilde{c}\in\mathcal{C}}\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \sum_{\tilde{t}\in\mathcal{T}}\xi_{\tilde{c}}^{\tilde{t}}(X)$ by definition of \tilde{X} , so \tilde{X} is a legitimate matching. Finally, $\xi_{\tilde{d}}^{\tilde{t}}(\tilde{X}) = \xi_{\tilde{d}}^{\tilde{t}}(X)$ for every \tilde{t} and \tilde{d} , so $\tilde{X} \in \mathcal{M}_1$.

matching such that

$$\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \begin{cases} \xi_{c_k}^{t_k}(X) + 1 & \text{ for } (\tilde{t}, \tilde{c}) = (t_k, c_k) \text{ for any } k \in \{i, i+1, \dots, j-1\}, \\ \xi_{c_k}^{t_{k+1}}(X) - 1 & \text{ for } (\tilde{t}, \tilde{c}) = (t_{k+1}, c_k^*) \text{ for any } k \in \{i, i+1, \dots, j-1\}, \\ \xi_{\tilde{c}}^{\tilde{t}}(X) & \text{ otherwise.} \end{cases}$$

We will show $\tilde{X} \in \mathcal{M}_1$. To do so, by construction of \tilde{X} , first note that $\sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) \leq \sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(X) + 1 \leq q_{\tilde{c}}$ for any $\tilde{c} \in \{c_i, \ldots, c_{j-1}\}$ such that $\sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(X) < q_{\tilde{c}}$. Next, by construction of \tilde{X} , $\sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(X) = q_{\tilde{c}}$ for every $\tilde{c} \in \{c_i, \ldots, c_{j-1}\}$ such that $\sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(X) = q_{\tilde{c}}$. Moreover, $\sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) \leq \sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(X) = q_{\tilde{c}}$ for every $\tilde{c} \in \{c_i^*, \ldots, c_{j-1}^*\}$. Finally, for every $\tilde{c} \in \mathcal{C} \setminus \{c_i, \ldots, c_{j-1}, c_i^*, \ldots, c_{j-1}^*\}$, $\sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \sum_{\tilde{t}\in\mathcal{T}} \xi_{\tilde{c}}^{\tilde{t}}(X) \leq q_{\tilde{c}}$. Thus, all school capacities are satisfied by \tilde{X} . Also by construction of \tilde{X} , for each $\tilde{d} \in \mathcal{D}$, $\sum_{\tilde{c}:d(\tilde{c})=\tilde{d}} \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \sum_{\tilde{c}:d(\tilde{c})=\tilde{d}} \xi_{\tilde{c}}^{\tilde{t}}(X) = k_{\tilde{d}}$, so \tilde{X} is rationed. Furthermore, for every $\tilde{c} \in \mathcal{C}$ and $\tilde{t} \in \mathcal{T}, \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) \leq \max\{\xi_{\tilde{c}}^{\tilde{t}}(X)\}$ by construction, so all type-specific ceilings are satisfied. Moreover, by construction of \tilde{X} , for each $\tilde{t} \in \mathcal{T}$, either $\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \xi_{\tilde{c}'}^{\tilde{t}}(\tilde{X}) + 1$, $\xi_{\tilde{c}''}^{\tilde{t}}(\tilde{X}) = \xi_{\tilde{c}'}^{\tilde{t}}(\tilde{X}) - 1$, and $\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) = \xi_{\tilde{c}}^{\tilde{t}}(X)$ for every $\tilde{c} \in \mathcal{C} \setminus \{\tilde{c}', \tilde{C}'\}$. Thus, $\tilde{t} \in \mathcal{T}$, $\xi_{\tilde{c}''}^{\tilde{t}}(\tilde{X}) = \xi_{\tilde{c}'}^{\tilde{t}}(\tilde{X}) = \xi_{\tilde{c}'}^{\tilde{t}}(\tilde{X})$ for every $\tilde{t} \in \mathcal{T}$. Therefore, \tilde{X} is legitimate.

By construction of \tilde{X} , either $\xi_{d'}^t(\tilde{X}) = \xi_{d'}^t(X)$ or $\xi_{d'}^t(\tilde{X}) = \xi_{d'}^t(X) - 1$. This implies that $\tilde{X} \in \mathcal{M}_1$. Furthermore, $\sum_{\tilde{t},\tilde{c}} |\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X}) - \xi_{\tilde{c}}^{\tilde{t}}(\hat{X})| < \sum_{\tilde{t},\tilde{c}} |\xi_{\tilde{c}}^{\tilde{t}}(X) - \xi_{\tilde{c}}^{\tilde{t}}(\hat{X})|$, since while creating the $\xi_{\tilde{c}}^{\tilde{t}}(\tilde{X})$ entries, we add 1 to some entries of X that satisfy $\xi_{\tilde{c}}^{\tilde{t}}(X) < \xi_{\tilde{c}}^{\tilde{t}}(\hat{X})$ and subtract 1 from some entries of X that satisfy $\xi_{\tilde{c}}^{\tilde{t}}(\hat{X}) > \xi_{\tilde{c}}^{\tilde{t}}(\hat{X})$. These lead to a contradiction to the assumption that $X \in \mathcal{M}_2$, which completes the proof.

Now we are ready to prove the theorem. The "if" part follows from Lemmas 2 and 3. Specifically, by Lemma 2, SPDA produces a legitimate matching. Therefore, by Lemma 3, we have $\hat{p}_d^t \leq \xi_d^t(X) \leq \hat{q}_d^t$ for every $t \in \mathcal{T}$ and $d \in \mathcal{D}$. For each school district d, hence, the maximum proportion of type-t students that can be admitted is \hat{q}_d^t/k_d and the minimum proportion of type t students that can be admitted is \hat{p}_d^t/k_d . Therefore, the ratio difference of type-t students in any two districts is at most $\max_{d\neq d'} \{\hat{q}_d^t/k_d - \hat{p}_{d'}^t/k_{d'}\}$. We conclude that the α -diversity policy is achieved when $\hat{q}_d^t/k_d - \hat{p}_{d'}^t/k_{d'} \leq \alpha$ for every t, d, and d' with $d \neq d'$.

The "only if" part of the theorem follows from Lemma 4. Suppose that $\hat{q}_d^t/k_d - \hat{p}_{d'}^t/k_{d'} > \alpha$ for some t, d, and d' with $d \neq d'$. From Lemma 4, we know the existence of a legitimate matching X such that $\xi_d^t(X) = \hat{q}_d^t$ and $\xi_{d'}^t(X) = \hat{p}_{d'}^t$. Consider a student preference profile where each student prefers her contract in X the most. Then, since the admissions rules are weakly acceptant, SPDA ends at the first step as all applications are accepted. Thus X is the outcome of SPDA and, therefore, the α -diversity policy is not satisfied.

Proof of Theorem 5. Suzuki et al. (2017) study a setting in which each student is initially endowed with a school and there are no constraints associated with student types, that is, when there is just one type. In that setting, they show that if the distribution is M-convex, then their mechanism, called TTC-M, satisfies the policy goal, constrained efficiency, individual rationality, and strategy-proofness. To adapt their result to our setting, consider the hypothetical matching problem that we have introduced before the definition of TTC in which each student is endowed with a school-type pair and each student has strict preferences over all school-type pairs. It is straightforward to verify that this hypothetical problem satisfies all the conditions assumed by Suzuki et al. (2017). In particular, M-convexity of $\Xi \cap \Xi^0$ holds by assumption. Therefore, TTC-M in this market satisfies the policy goal, constrained efficiency, individual rationality, and strategy-proofness.

We note that the outcome of our TTC is isomorphic to the outcome of TTC-M in the hypothetical problem in the following sense. Student *s* is allocated to contract (s, c) under preference profile $P = (P_s)_{s \in S}$ at the outcome of the TTC mechanism if, and only if, student *s* is allocated to the school-type pair (c, t) under preference profile $\tilde{P} = (\tilde{P}_s)_{s \in S}$ at TTC-M in the hypothetical problem. The rest of the proof is devoted to showing that our TTC satisfies the desired properties in the original problem.

The result that TTC satisfies the policy goal follows from the result in Suzuki et al. (2017) that the distribution corresponding to the TTC-M outcome is in $\Xi \cap \Xi^0$.

To show constrained efficiency, let X be the outcome of TTC and, for each student $s \in S$, let (s, c_s) be the contract associated with student s at matching X. Suppose, for contradiction, that there exists a feasible matching X' with $\xi(X') \in \Xi$ that Pareto dominates matching X. Denoting $X'_s = (s, c'_s)$ for each student $s \in S$, this implies $(s, c'_s) R_s (s, c_s)$ for every student $s \in S$, with at least one relation being strict. Then, by the construction of preferences \tilde{R}_s in the hypothetical problem, we have $(c'_s, \tau(s)) \tilde{R}_s (c_s, \tau(s))$ for every student $s \in S$, with at least one relation being strict. Moreover, because matching X' is feasible in the original problem, $Y' = \{(c'_s, \tau(s)) | (s, c'_s) \in X'\}$ is feasible in the hypothetical problem, and $Y = \{(c_s, \tau(s)) | (s, c_s) \in X\}$ is the result of TTC-M. This is a contradiction to the result in Suzuki et al. (2017) that TTC-M is constrained efficient.

To show individual rationality, let matching X be the outcome of TTC and, for each student $s \in S$, let $X_s = (s, c_s)$ be the contract associated with student s at matching X. Additionally, let $Y = \{(c_s, \tau(s)) | (s, c_s) \in X\}$ be the result of TTC-M in the hypothetical problem. Suzuki et al. (2017) establish that TTC-M is individually rational, so $(c_s, \tau(s)) \tilde{R}_s$ $(c_0(s), \tau(s))$ for every $s \in S$, where $c_0(s)$ denotes the initial school of student s. By the construction of \tilde{R}_s , this relation implies $(s, c_s) R_s (s, c_0(s))$ for every student $s \in S$, which means X is individually rational in the original problem. To show strategy-proofness, in the original problem, let *s* be a student, *t* her type, P_{-s} the preference profile of students other than student *s*, P_s the true preference of student *s*, and P'_s a misreported preference of student *s*. Furthermore, let *c* and *c'* be schools assigned to student *s* under (P_s, P_{-s}) and (P'_s, P_{-s}) for TTC, respectively. Note that the previous argument establishes that, in the hypothetical problem, student *s* is allocated to (c, t) and (c', t) under $(\tilde{P}_s, \tilde{P}_{-s})$ and $(\tilde{P}'_s, \tilde{P}'_{-s})$, respectively. Because TTC-M is strategy-proof, it follows that $(c, t) \ \tilde{P}_s \ (c', t)$ or c = c'. By the construction of \tilde{P}_s , this relation implies $(s, c) \ P_s \ (s, c')$ or (s, c) = (s, c'), establishing strategy-proofness of TTC in the original problem.

Proof of Corollary 1. Assume that $\sum_t \xi_c^t \leq q_c$ for every $\xi \in \Xi$ and $c \in C$. Under this presumption, we show that when the policy goal Ξ is M-convex, so is $\Xi \cap \Xi^0$. Then the result follows immediately from Theorem 5 because the initial matching satisfies Ξ .

Suppose that $\xi, \tilde{\xi} \in \Xi \cap \Xi^0$ such that $\xi_c^t > \tilde{\xi}_c^t$ for some school c and type t. Since Ξ is M-convex and $\xi, \tilde{\xi} \in \Xi$, there exist school c' and type t' with $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ such that $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi$ and $\tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi$. We need to show that $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi^0$ and $\tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi^0$ as well.

Because $\xi \in \Xi^0$, the number of students assigned in ξ is $\sum_d k_d$. Therefore, the number of students assigned in $\xi - \chi_{c,t} + \chi_{c',t'}$ is also $\sum_d k_d$. Furthermore, $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi$ implies that the number of students in a school at $\xi - \chi_{c,t} + \chi_{c',t'}$ is less than or equal to the capacity of the school. These two results imply that $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi^0$. \Box

Proof of Lemma 1. Note that $\Xi(f, \lambda) \cap \Xi^0 = \{\xi \in \Xi_0 | f(\xi) \ge \lambda\}.$

The "if" direction: Suppose that $\xi \in \Xi(f, \lambda) \cap \Xi^0$ and $\tilde{\xi} \in \Xi(f, \lambda) \cap \Xi^0$ are distinct. Therefore, $f(\xi), f(\tilde{\xi}) \ge \lambda$. By assumption, there exist (c, t) and (c', t') with $\xi_c^t > \tilde{\xi}_c^t$ and $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ such that

$$\min\{f(\xi - \chi_{c,t} + \chi_{c',t'}), f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'})\} \ge \min\{f(\xi), f(\tilde{\xi})\}.$$

This implies $f(\xi - \chi_{c,t} + \chi_{c',t'})$, $f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'}) \ge \lambda$. Furthermore, $\xi - \chi_{c,t} + \chi_{c',t'}, \tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi^0$ since the sum of coordinates is equal to $\sum_d k_d$ and no school is assigned more students than its capacity. Therefore, $\xi - \chi_{c,t} + \chi_{c',t'}, \tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi(f,\lambda) \cap \Xi^0$, so, by Theorem 4.3. of Murota (2003), $\Xi(f,\lambda) \cap \Xi^0$ is M-convex.

The "only if" direction: Suppose that the function f is not pseudo M-concave, so that there exist distinct $\xi, \tilde{\xi} \in \Xi_0$ such that for all (c, t) and (c', t') with $\xi_c^t > \tilde{\xi}_c^t$ and $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ we have

$$\min\{f(\xi - \chi_{c,t} + \chi_{c',t'}), f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'})\} < \min\{f(\xi), f(\tilde{\xi})\}.$$

Let $\lambda \equiv \min\{f(\xi), f(\tilde{\xi})\}$. The above condition implies that $\Xi(f, \lambda) \cap \Xi^0$ is not M-convex. \Box

Proof of Theorem 7. Let $f(\xi) = 1$ when $\xi \in \Xi \cap \Xi^0$ and $f(\xi) = 0$ otherwise.

First we show that f is pseudo M-concave. Take two distinct $\xi, \tilde{\xi} \in \Xi_0$. If $\min\{f(\xi), f(\tilde{\xi})\} = 0$, then $\min\{f(\xi - \chi_{c,t} + \chi_{c',t'}), f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'})\} \ge 0$ for every (c,t) and (c',t'), so the desired inequality holds. Suppose that $f(\xi) = f(\tilde{\xi}) = 1$. By the construction of f, we have $\xi, \tilde{\xi} \in \Xi \cap \Xi^0$. Since $\Xi \cap \Xi^0$ is M-convex, there exist (c,t) and (c',t') such that $\xi - \chi_{c,t} + \chi_{c',t'}, \tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi \cap \Xi^0$. By the construction of f, $f(\xi - \chi_{c,t} + \chi_{c',t'}) = f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'}) = 1$. Therefore, the desired inequality also holds for this case, so f is pseudo M-concave.

Next we show that $\Xi(f,\lambda) \cap \Xi^0 = \Xi \cap \Xi^0$ for $\lambda = 1$. For any $\xi \in \Xi(f,1) \cap \Xi^0$, $f(\xi) = 1$, which implies that $\xi \in \Xi \cap \Xi^0$ by the construction of f. Therefore, $\Xi(f,1) \cap \Xi^0 \subseteq \Xi \cap \Xi^0$. Now, let $\xi \in \Xi \cap \Xi^0$. Then, by the construction of f, $f(\xi) = 1$, so $\xi \in \Xi(f,1) \cap \Xi^0$. Therefore, $\Xi \cap \Xi^0 \subseteq \Xi(f,1) \cap \Xi^0$. We conclude that $\Xi(f,1) \cap \Xi^0 = \Xi \cap \Xi^0$.

Proof of Corollary 2. Suppose that the diversity policy Ξ sets a floor and ceiling for each type at each school. We show that $\Xi \cap \Xi^0$ is an M-convex set. The set $\Xi \cap \Xi^0$ can be represented as $\{\xi | \forall c, t \ q_c^t \ge \xi_c^t \ge p_c^t, \ \forall c \ q_c \ge \sum_t \xi_c^t \text{ and } \sum_{c,t} \xi_c^t = \sum_d k_d \}.$

Suppose that there exist $\xi, \tilde{\xi} \in \Xi \cap \Xi^0$ such that $\xi_c^t > \tilde{\xi}_c^t$. To show M-convexity, we need to find school c' and type t' with $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ such that (1) $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi \cap \Xi^0$ and (2) $\tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi \cap \Xi^0$. To show both conditions, we look at two possible cases depending on whether c' = c or not.

Case 1: First consider the case in which there exists type t' such that $\xi_c^{t'} < \tilde{\xi}_c^{t'}$. We prove (1) that $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi \cap \Xi^0$. Since $\xi - \chi_{c,t} + \chi_{c,t'}$ assigns the same total number of students at school c as ξ , the capacity constraint at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c,t'}$. Furthermore, the number of students assigned to any school in $\xi - \chi_{c,t} + \chi_{c,t'}$ is the same as ξ , which means that all students are assigned. Next, because $\xi_c^t - 1 \ge \tilde{\xi}_c^t \ge p_c^t$ (the former inequality comes from the assumption $\xi_c^t > \tilde{\xi}_c^t$, and the latter comes from the fact that $\tilde{\xi} \in \Xi$), the floor for type t at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c,t'}$.

The floor for type t' at school c is satisfied for $\xi - \chi_{c,t} + \chi_{c,t'}$ because $\xi_c^{t'} + 1 \ge \xi_c^{t'} \ge p_c^{t'}$ (the former inequality is obvious, and the latter comes from the fact that $\xi \in \Xi$). Similarly, the ceiling for type t' at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c,t'}$ because $q_c^t \ge \tilde{\xi}_c^{t'} \ge \xi_c^{t'} + 1$.

No other coefficients changed between ξ and $\xi - \chi_{c,t} + \chi_{c,t'}$, so all other constraints are satisfied at the latter distribution. Therefore, (1) is satisfied.

The proof that (1) is satisfied follows from the facts that $\xi_c^t > \tilde{\xi}_c^t$ and $\xi_c^{t'} < \tilde{\xi}_c^{t'}$. By changing the roles of t with t' and ξ with $\tilde{\xi}$ in the preceding argument, we get the implication of (1) that $\tilde{\xi} - \chi_{c,t'} + \chi_{c,t} \in \Xi \cap \Xi^0$. But this is exactly (2).

Case 2: Second, consider the case in which there exists no type t' such that $\xi_c^{t'} < \tilde{\xi}_c^{t'}$. Then, $\xi_c^{t'} \ge \tilde{\xi}_c^{t'}$ for every $t' \ne t$. In particular, the total number of students assigned to school c at ξ is strictly larger than at $\tilde{\xi}$. Because everyone is matched with some school at ξ and $\tilde{\xi}$ by assumption, it is implied that, without loss of generality, there exists school $c' \ne c$ such that the total number of students matched with c' is strictly larger at $\tilde{\xi}$ than at ξ . In addition, there exists type t' such that $\tilde{\xi}_{c'}^{t'} > \xi_{c'}^{t'}$.

Now we proceed to show condition (1) for this case. To do so, we first note that $\xi - \chi_{c,t} + \chi_{c',t'}$ assigns the same number of students as in ξ , so all students are assigned in $\xi - \chi_{c,t} + \chi_{c',t'}$. Furthermore, it assigns a smaller number of students at school c than ξ , so the capacity constraint at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$. Likewise, $\xi - \chi_{c,t} + \chi_{c',t'}$ assigns a weakly smaller number of students at c' than $\tilde{\xi}$ does, so the capacity constraint at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$.

Next we check that the floor for type t and ceiling for type t' at school c are satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$. Because $\xi_c^t - 1 \ge \tilde{\xi}_c^t \ge p_c^t$ (the first inequality follows from the assumption $\xi_c^t > \tilde{\xi}_c^t$ and the second from the fact that $\tilde{\xi} \in \Xi$), the floor for type t at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$. Since $q_c^t \ge \xi_c^t > \xi_c^t - 1$ (because $\xi \in \Xi$), the ceiling for type t at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$.

Now we check that the floor for type t' and ceiling for type t at school c' are satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$. For type t' at school c', we have $\xi_{c'}^{t'} + 1 > \xi_{c'}^{t'} \ge p_{c'}^{t'}$ (the first inequality is obvious and the second follows from the fact that $\xi \in \Xi$), so the floor for type t' at school c' is satisfied for $\xi - \chi_{c,t} + \chi_{c',t'}$. Furthermore, we have $q_{c'}^{t'} \ge \tilde{\xi}_{c'}^{t'} \ge \xi_{c'}^{t'} + 1$ (the first inequality follows from the fact that $\tilde{\xi} \in \Xi$ and the second one follows from $\tilde{\xi}_{c'}^{t'} > \xi_{c'}^{t'}$), so the ceiling for type t' at school c' is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$.

No other coefficients changed between ξ and $\xi - \chi_{c,t} + \chi_{c',t'}$, so all other constraints are satisfied at the latter distribution. Therefore, (1) is satisfied.

The proof that (1) is satisfied follows from the facts that $\xi_c^t > \tilde{\xi}_{c'}^t \tilde{\xi}_{c'}^{t'} > \xi_{c'}^{t'}$, there are more students assigned to school c at ξ than $\tilde{\xi}$, and there are more students assigned to school c' at $\tilde{\xi}$ than ξ . If we change the roles of ξ with $\tilde{\xi}$, c with c', and t with t', then (1) would imply $\tilde{\xi} - \chi_{c',t'} + \chi_{c,t} \in \Xi \cap \Xi^0$. But this is exactly (2), so we are done. Therefore, $\Xi \cap \Xi^0$ is an M-convex set.

The desired conclusion then follows from Theorem 5.

Proof of Corollary 3. We show that f is pseudo M-concave. Let $\xi, \tilde{\xi} \in \Xi^0$ be distinct. Then $U \equiv \{(c,t) | \xi_t^c > \tilde{\xi}_t^c\}$ is a nonempty set. Partition this set into three subsets: $U_1 \equiv \{(c,t) | \hat{\xi}_t^c \ge \xi_t^c > \tilde{\xi}_t^c\}, U_2 \equiv \{(c,t) | \xi_t^c > \hat{\xi}_t^c > \tilde{\xi}_t^c\}, \text{ and } U_3 \equiv \{(c,t) | \xi_t^c > \tilde{\xi}_t^c \ge \hat{\xi}_t^c\}$. Likewise $V \equiv \{(c',t') | \tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'}\}$ is a nonempty set that can be partitioned into three subsets: $V_1 \equiv \{(c',t') | \hat{\xi}_{t'}^{c'} \ge \tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'} \}, V_2 \equiv \{(c',t') | \tilde{\xi}_{t'}^{c'} > \hat{\xi}_{t'}^{c'} > \xi_{t'}^{c'} \}, \text{ and } V_3 \equiv \{(c',t') | \tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'} \ge \hat{\xi}_{t'}^{c'} \}.$ We consider several cases.

Case 1: U_2 is nonempty. There exists (c, t) such that $\xi_t^c > \hat{\xi}_t^c > \tilde{\xi}_t^c$. Since $\xi, \tilde{\xi} \in \Xi_0$, there exists (c', t') such that $\xi_{t'}^{c'} < \tilde{\xi}_{t'}^{c'}$. In this case, $f(\xi - \chi_{c,t} + \chi_{c',t'}) \ge f(\xi)$ and $f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'}) \ge f(\tilde{\xi})$ by definition of f. Therefore,

$$\min\{f(\xi - \chi_{c,t} + \chi_{c',t'}), f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'})\} \ge \min\{f(\xi), f(\tilde{\xi})\}$$

Case 2: V_2 is nonempty. There exists (c', t') such that $\tilde{\xi}_{t'}^{c'} > \hat{\xi}_{t'}^{c'} > \xi_{t'}^{c'}$. The proof of this case is similar to the proof of Case 1.

Case 3: U_1 and V_1 are nonempty. There exist (c,t) and (c',t') such that $\hat{\xi}_t^c \ge \xi_t^c > \tilde{\xi}_t^c$ and $\hat{\xi}_{t'}^{c'} \ge \tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'}$. In this case, $f(\xi - \chi_{c,t} + \chi_{c',t'}) = f(\xi)$ and $f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'}) = f(\tilde{\xi})$ by definition of f. Then,

$$\min\{f(\xi - \chi_{c,t} + \chi_{c',t'}), f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'})\} = \min\{f(\xi), f(\tilde{\xi})\}.$$

Case 4: U_3 and V_3 are nonempty. There exist (c, t) and (c', t') such that $\xi_t^c > \tilde{\xi}_t^c \ge \hat{\xi}_t^c$ and $\tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'} \ge \hat{\xi}_{t'}^{c'}$. The proof is similar to the proof of Case 3.

Case 5: $U = U_1$ and $V = V_3$. For every (c, t) such that $\xi_t^c > \tilde{\xi}_t^c$, we have $\hat{\xi}_t^c \ge \xi_t^c > \tilde{\xi}_t^c$, and for every (c', t') such that $\tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'} > \xi_{t'}^{c'} \ge \xi_{t'}^{c'}$. In this case, $f(\tilde{\xi}) + 2 \le f(\xi)$, so $\min\{f(\tilde{\xi}), f(\xi)\} = f(\tilde{\xi})$. Furthermore, for any choice of (c, t) and (c', t') such that $\xi_c^t > \tilde{\xi}_c^t$ and $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$, we have $f(\xi - \chi_{c,t} + \chi_{c',t'}) = f(\xi) - 2$ and $f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'}) = f(\tilde{\xi}) + 2$.

Since $f(\xi) - 2 \ge f(\tilde{\xi})$ and $f(\tilde{\xi}) + 2 > f(\tilde{\xi})$, we get the desired conclusion that

$$\min\{f(\xi - \chi_{c,t} + \chi_{c',t'}), f(\tilde{\xi} + \chi_{c,t} - \chi_{c',t'})\} \ge f(\tilde{\xi}) = \min\{f(\xi), f(\tilde{\xi})\}.$$

Case 6: $U = U_3$ and $V = V_1$. For every (c, t) such that $\xi_t^c > \tilde{\xi}_t^c$, we have $\xi_t^c > \tilde{\xi}_t^c \ge \hat{\xi}_t^c$, and for every (c', t') such that $\tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'}$, we have $\hat{\xi}_{t'}^{c'} \ge \tilde{\xi}_{t'}^{c'} > \xi_{t'}^{c'}$. The proof is similar to the proof of Case 5.

We have considered all the possible cases: If U_2 or V_2 are nonempty, then we are done by Cases 1 and 2, respectively. Suppose that they are both empty. Therefore, $U = U_1 \cup U_3$ and $V = V_1 \cup V_3$ are both nonempty. If U_1 and V_1 are nonempty, then we are done by Case 3. If U_3 and V_3 are nonempty, then we are done by Case 4. If U_1 and V_3 are nonempty and one of U_3 or V_1 is nonempty, then we are done by Cases 3 or 4. Otherwise, if U_3 and V_1 are empty when U_1 and V_3 are nonempty, then $U = U_1$ and $V = V_3$, which is covered by Case 5. If U_3 and V_1 are nonempty and one of U_1 or V_3 is nonempty, then we are done by Cases 3 or 4. Otherwise, if U_1 and V_3 are empty when U_3 and V_1 are nonempty, then $U = U_3$ and $V = V_1$, which is covered by Case 6.

We conclude that f is pseudo M-concave since in all possible cases we derive the desired inequality. Then the proof follows from Theorem 6.

Proof of Corollary 4. Let the balanced-exchange policy be denoted by Ξ . We show that $\Xi \cap \Xi^0 = \{\xi | \forall d \sum_t \xi_d^t = k_d \text{ and } \forall c q_c \ge \sum_t \xi_c^t \}$ is M-convex.

Suppose that there exist $\xi, \tilde{\xi} \in \Xi \cap \Xi^0$ such that $\xi_c^t > \tilde{\xi}_c^t$. To show M-convexity, we need to find school c' and type t' with $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ such that (1) $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi \cap \Xi^0$ and (2) $\tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi \cap \Xi^0$.

If there exists t' such that $\tilde{\xi}_c^{t'} > \xi_c^{t'}$, then the number of students in each district and each school are the same in ξ with $\xi - \chi_{c,t} + \chi_{c,t'}$ and $\tilde{\xi}$ with $\tilde{\xi} + \chi_{c,t} - \chi_{c,t'}$, so both (1) and (2) are satisfied.

Otherwise, suppose that, for every type $t' \neq t$, $\tilde{\xi}_c^{t'} \leq \xi_c^{t'}$. Therefore, there are more students assigned to school c at ξ than $\tilde{\xi}$. Since the number of students assigned to district $d \equiv d(c)$ in ξ and $\tilde{\xi}$ are the same, there exists another school c' in district d such that c' has more students in $\tilde{\xi}$ than ξ . Furthermore, there exists type t' such that $\tilde{\xi}_{c'}^{t'} > \xi_{c'}^{t'}$.

We first show (1). Since both schools c and c' are in district d, the number of students assigned to district d is the same at ξ and $\xi - \chi_{c,t} + \chi_{c',t'}$. Therefore, the number of students assigned to district d at $\xi - \chi_{c,t} + \chi_{c',t'}$ is k_d .

Next we check the school capacity constraints. The number of students assigned to school c at $\xi - \chi_{c,t} + \chi_{c',t'}$ is one less than the corresponding number at ξ , so the capacity constraint of school c at $\xi - \chi_{c,t} + \chi_{c',t'}$ is satisfied. Furthermore, the number of students assigned to school c' at $\xi - \chi_{c,t} + \chi_{c',t'}$ is weakly smaller than the corresponding number at $\tilde{\xi}$. Therefore, the capacity constraint of school c' at $\xi - \chi_{c,t} + \chi_{c',t'}$ is also satisfied.

Since all the other coefficients are the same at ξ and $\xi - \chi_{c,t} + \chi_{c',t'}$, (1) holds.

Note that the above argument relies on the facts $\xi_c^t > \tilde{\xi}_c^t, \xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$, and d(c) = d(c'). If we switch the roles of c with c' and ξ with $\tilde{\xi}$, the implication of (1) is $\tilde{\xi} - \chi_{c',t'} + \chi_{c,t} \in \Xi \cap \Xi^0$, which is exactly (2). Therefore, Ξ is M-convex.

The result then follows from Theorem 5 because $\Xi \cap \Xi^0$ is M-convex and the initial matching trivially satisfies the balanced-exchange policy.

Proof of Corollary 5. We first show that the set of distributions $\Xi \cap \Xi^0 = \{\xi | \forall c, t q_c^t \ge \xi_c^t \ge p_c^t, \forall c q_c \ge \sum_t \xi_c^t \text{ and } \forall d \sum_t \xi_d^t = k_d\}$ is an M-convex set.

Suppose that there exist $\xi, \tilde{\xi} \in \Xi \cap \Xi^0$ such that $\xi_c^t > \tilde{\xi}_c^t$. To show M-convexity, we need to find school c' and type t' with $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$ such that (1) $\xi - \chi_{c,t} + \chi_{c',t'} \in \Xi \cap \Xi^0$ and (2) $\tilde{\xi} + \chi_{c,t} - \chi_{c',t'} \in \Xi \cap \Xi^0$. Let $d \equiv d(c)$. To show both conditions, we look at two possible cases depending on whether c' = c or not.

Case 1: First consider the case when there exists type t' such that $\xi_c^{t'} < \tilde{\xi}_c^{t'}$. We prove (1) that $\xi - \chi_{c,t} + \chi_{c,t'} \in \Xi \cap \Xi^0$. Since $\xi - \chi_{c,t} + \chi_{c,t'}$ assigns the same total number of students at school c as ξ , the capacity constraint at school c at $\xi - \chi_{c,t} + \chi_{c,t'}$ is satisfied. Furthermore, the number of students assigned to any district at $\xi - \chi_{c,t} + \chi_{c,t'}$ is the same

as ξ , which means that the number of students in every district is equal to the number of students who are from there. Next, because $\xi_c^t - 1 \ge \tilde{\xi}_c^t \ge p_c^t$ (the former inequality comes from the assumption $\xi_c^t > \tilde{\xi}_{c'}^t$ and the latter comes from the fact $\tilde{\xi} \in \Xi \cap \Xi^0$), the floor for type t at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c,t'}$. Next, the facts that $\xi \in \Xi \cap \Xi^0$ and $\xi_c^t > \tilde{\xi}_c^t$ implies $q_c^t \ge \xi_c^t \ge \tilde{\xi}_c^t + 1$. Therefore, the ceiling for type t at school c at $\xi - \chi_{c,t} + \chi_{c,t'}$ is satisfied.

The floor for type t' at school c is satisfied for $\xi - \chi_{c,t} + \chi_{c,t'}$ because $\xi_c^{t'} + 1 \ge \xi_c^{t'} \ge p_c^{t'}$ (the former inequality is obvious, and the latter comes from the fact $\xi \in \Xi \cap \Xi^0$). Similarly, the ceiling for type t' at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c,t'}$ because $q_c^t \ge \tilde{\xi}_c^{t'} \ge \xi_c^{t'} + 1$.

No other coefficients changed between ξ and $\xi - \chi_{c,t} + \chi_{c,t'}$, so all other constraints are satisfied at the latter distribution. Therefore, (1) is satisfied.

The proof that (1) is satisfied follows from the facts that $\xi_c^t > \tilde{\xi}_c^t$ and $\xi_c^{t'} < \tilde{\xi}_c^{t'}$. By changing the roles of t with t' and ξ with $\tilde{\xi}$ in the preceding argument, we get the implication of (1) that $\tilde{\xi} - \chi_{c,t'} + \chi_{c,t} \in \Xi \cap \Xi^0$. But this is exactly (2).

Case 2: In this case, $c' \neq c$ for every (c', t') such that $\xi_{c'}^{t'} < \tilde{\xi}_{c'}^{t'}$. Then, $\xi_c^{t'} \geq \tilde{\xi}_c^{t'}$ for every $t' \neq t$. In particular, the total number of students assigned to school c at ξ is strictly larger than at $\tilde{\xi}$. Because the number of students in district d are the same at ξ and $\tilde{\xi}$, there exist school c' in district d such that the total number of students matched with c' is strictly larger at $\tilde{\xi}$ than at ξ . In addition, there exists type t' such that $\tilde{\xi}_{c'}^{t'} > \xi_{c'}^{t'}$.

Now we proceed to show condition (1) for this case. To do so, we first note that $\xi - \chi_{c,t} + \chi_{c',t'}$ assigns the same number of students to each district as in ξ , so the number of students assigned to each district d is k_d at $\xi - \chi_{c,t} + \chi_{c',t'}$. Furthermore, it assigns a smaller number of students at school c than ξ , so the capacity constraint at school c at $\xi - \chi_{c,t} + \chi_{c',t'}$ is satisfied. Likewise, $\xi - \chi_{c,t} + \chi_{c',t'}$ assigns a weakly smaller number of students at c' than $\tilde{\xi}$ does, so the capacity constraint at school c' at $\xi - \chi_{c,t} + \chi_{c',t'}$ is satisfied.

Next we check that the type-specific floors at schools are satisfied. Because $\xi_c^t - 1 \ge \tilde{\xi}_c^t \ge p_c^t$ (the first inequality follows from the assumption $\xi_c^t > \tilde{\xi}_c^t$ and the second from the fact that $\tilde{\xi} \in \Xi \cap \Xi^0$), the floor for type t at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$. For type t' at school c', we have $\xi_{c'}^{t'} + 1 \ge \xi_{c'}^{t'} \ge p_{c'}^{t'}$ (the first inequality is obvious and the second follows from the fact that $\xi \in \Xi \cap \Xi^0$), so the floor for type t' at school c' is satisfied for $\xi - \chi_{c,t} + \chi_{c',t'}$.

Now we check that the type-specific ceilings at schools are satisfied. Since $q_c^t \ge \xi_c^t > \xi_c^t - 1$ (the first inequality follows from the fact that $\xi \in \Xi \cap \Xi^0$ and the second inequality is obvious), the ceiling for type t at school c is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$. For type t' at school c', we have $q_{c'}^{t'} \ge \tilde{\xi}_{c'}^{t'} \ge \xi_{c'}^{t'} + 1$ (the first inequality follows from the fact that $\tilde{\xi} \in \Xi \cap \Xi^0$ and the second one follows from $\tilde{\xi}_{c'}^{t'} > \xi_{c'}^{t'}$), so the ceiling for type t' at school c' is satisfied at $\xi - \chi_{c,t} + \chi_{c',t'}$.

No other coefficients changed between ξ and $\xi - \chi_{c,t} + \chi_{c',t'}$, so all other constraints are satisfied at the latter distribution.

The proof that (1) is satisfied follows from the facts that d(c) = d(c'), $\xi_c^t > \tilde{\xi}_c^t$, $\tilde{\xi}_{c'}^{t'} > \xi_{c'}^{t'}$, there are more students assigned to school c at ξ than $\tilde{\xi}$, and there are more students assigned to school c' at $\tilde{\xi}$ than ξ . If we change the roles of ξ with $\tilde{\xi}$, c with c', and t with t', then (1) would imply $\tilde{\xi} - \chi_{c',t'} + \chi_{c,t} \in \Xi \cap \Xi^0$. But this is exactly (2), so we are done.

The result then follows from Theorem 5 because $\Xi \cap \Xi^0$ is M-convex.

Proof of Theorem 8. Suppose that district admissions rules favor their own students. Fix a student preference profile. Recall that under interdistrict school choice, students are assigned to schools by SPDA, where each student ranks all contracts associated with her and each district d has the admissions rule Ch_d . Under intradistrict school choice, students are assigned to schools by SPDA where students only rank the contracts associated with their home districts and each district d has the admissions rule Ch_d . We first show that the intradistrict SPDA outcome can be produced by SPDA when all districts participate simultaneously and students rank all contracts, including the ones associated with the other districts, by modifying admissions rules for the districts. Let $Ch'_d(X) \equiv Ch_d(\{x \in X | d(s(x)) = d\})$ be the modified admissions rule.

In SPDA, if district admissions rules have completions that satisfy path independence, then SPDA outcomes are the same under the completions and the original admissions rules because in SPDA a district always considers a set of proposals which is feasible for students. Furthermore, SPDA does not depend on the order of proposals when district admissions rules are path independent. As a result, SPDA does not depend on the order of proposals when district admissions rules have completions that satisfy path independence. Therefore, the intradistrict SPDA outcome can be produced by SPDA when all districts participate simultaneously and students rank all contracts including the ones associated with the other districts and each district *d* has the admissions rule Ch'_d . The reason behind this is that when each district *d* has admissions rule Ch'_d , a student is not admitted to a school district other than her home district. Furthermore, because Ch_d favors own students, the set of chosen students under Ch'_d is the same as that under Ch_d for any set of contracts of the form $\{x \in X | d(s(x)) = d\}$ for any set *X*.

We next show that Ch'_d has a path-independent completion. By assumption, for every district d, there exists a path-independent completion \widetilde{Ch}_d of Ch_d . Let $\widetilde{Ch}'_d(X) \equiv \widetilde{Ch}_d(\{x \in X | d(s(x)) = d\})$. We show that \widetilde{Ch}'_d is a path-independent completion of Ch'_d . To show that $\widetilde{Ch}'_d(X)$ is a completion, consider a set X such that $\widetilde{Ch}'_d(X)$ is feasible for students. Let $X^* \equiv \{x \in X | d(s(x)) = d\}$. Then we have the following:

$$Ch_d(X^*) = Ch_d(X^*) = Ch_d(X^*) = Ch_d(X^*),$$

where the first equality follows from the definition of \widetilde{Ch}'_d , the second equality follows from the fact that \widetilde{Ch}_d is a completion of Ch_d , and the third equality follows from the definition of Ch'_d . Furthermore, because $\widetilde{Ch}'_d(X) = \widetilde{Ch}'_d(X^*)$ and $Ch'_d(X^*) = Ch'_d(X)$, we get $\widetilde{Ch}'_d(X) = Ch'_d(X)$. Therefore, \widetilde{Ch}'_d is a completion of Ch'_d .

To show that \widetilde{Ch}'_d is path independent, consider two sets of contracts X and Y. Let $X^* \equiv \{x \in X | d(s(x)) = d\}$ and $Y^* \equiv \{x \in Y | d(s(x)) = d\}$. Then we have the following:

$$\widetilde{Ch}'_d(X \cup \widetilde{Ch}'_d(Y)) = \widetilde{Ch}'_d(X \cup \widetilde{Ch}_d(Y^*))$$
$$= \widetilde{Ch}_d(X^* \cup \widetilde{Ch}_d(Y^*))$$
$$= \widetilde{Ch}_d(X^* \cup Y^*)$$
$$= \widetilde{Ch}'_d(X \cup Y),$$

where the first and second equalities follow from the definition of \widetilde{Ch}'_d , the third equality follows from path independence of \widetilde{Ch}_d , and the last equality follows from the definition of \widetilde{Ch}'_d . Therefore, \widetilde{Ch}'_d is path independent.

Because Ch_d favors own students, we have $Ch_d(X) \supseteq Ch'_d(X)$ for every X that is feasible for students. Furthermore, for any such X, $\widetilde{Ch}_d(X) = Ch_d(X)$ and $\widetilde{Ch}'_d(X) = Ch'_d(X)$ because \widetilde{Ch}_d is a completion of Ch_d and \widetilde{Ch}'_d is a completion of Ch'_d , respectively. Therefore, for any X that is feasible for students, $\widetilde{Ch}_d(X) \supseteq \widetilde{Ch}'_d(X)$. We use this result to show the following lemma.

Lemma 5. Every student weakly prefers the interdistrict SPDA outcome under $(\widetilde{Ch}_d)_{d\in\mathcal{D}}$ to the interdistrict SPDA outcome under $(\widetilde{Ch}'_d)_{d\in\mathcal{D}}$.

Proof. Let μ be the interdistrict SPDA outcome under $(\widetilde{Ch}_d)_{d\in\mathcal{D}}$ and μ' be the interdistrict SPDA outcome under $(\widetilde{Ch}'_d)_{d\in\mathcal{D}}$. If μ' is stable under $(\widetilde{Ch}_d)_{d\in\mathcal{D}}$, then the conclusion follows from the result that μ is the student-optimal stable matching under $(\widetilde{Ch}_d)_{d\in\mathcal{D}}$ because each \widetilde{Ch}_d is path independent (Chambers and Yenmez, 2017).

Suppose that μ' is not stable under $(\widetilde{Ch}_d)_{d\in\mathcal{D}}$. Since μ' is stable under $(\widetilde{Ch}'_d)_{d\in\mathcal{D}}$, $\widetilde{Ch}'_d(\mu'_d) = \mu'_d$ for every district d. Furthermore, μ'_d is feasible for students, so $\widetilde{Ch}_d(\mu'_d) \supseteq \widetilde{Ch}'_d(\mu'_d) = \mu'_d$. By definition of admissions rules, $\mu'_d \supseteq \widetilde{Ch}_d(\mu'_d)$, so $\widetilde{Ch}_d(\mu'_d) = \mu'_d$. As a result, there must exist a blocking contract for matching μ' so that it is not stable under $(\widetilde{Ch}_d)_{d\in\mathcal{D}}$. Whenever there exists a blocking pair, we consider the following algorithm to improve student welfare. Let d_1 be the district associated with a blocking contract. Set $\mu^0 \equiv \mu'$.

Step n (n \ge 1): Consider the following set of contracts associated with district d_n for which there exists an associated blocking contract: $X_{d_n}^n \equiv \{x = (s, d_n, c) | x P_s \mu_s^{n-1}\}$.

District d_n accepts $Ch_{d_n}(\mu_d^{n-1} \cup X_{d_n}^n)$ and rejects the rest of the contracts. Let $\mu_{d_n}^n \equiv \widetilde{Ch}_{d_n}(\mu_{d_n}^{n-1} \cup X_{d_n}^n)$ and $\mu_d^n \equiv \mu_d^{n-1} \setminus Y^n$ where $Y^n \equiv \{x \in \mu^{n-1} | \exists y \in \mu_{d_n}^n \text{ s.t. } s(x) = s(y)\}$ for $d \neq d_n$. If there are no blocking contracts for matching μ^n under $(\widetilde{Ch}_d)_{d \in \mathcal{D}}$, then stop and return μ^n , otherwise go to Step n + 1.

We show that district d_n does not reject any contract in $\mu_{d_n}^{n-1}$ by mathematical induction on *n*, i.e., $\mu_{d_n}^n \supseteq \mu_{d_n}^{n-1}$ for every $n \ge 1$. Consider the base case for n = 1. Recall that $\mu_{d_1}^1 =$ $\widetilde{Ch}_{d_1}(\mu^0_{d_1} \cup X^1_{d_1}) = \widetilde{Ch}_{d_1}(\mu'_{d_1} \cup X^1_{d_1})$. By construction, $\mu^1_{d_1}$ is a feasible matching. We claim that $\mu'_{d_1} \cup \mu^1_{d_1}$ is feasible for students. Suppose, for contradiction, that it is not feasible for students. Then there exists a student *s* who has one contract in μ'_{d_1} and one in $\mu^1_{d_1} \setminus \mu'_{d_1}$. Call the latter contract z. By construction, $z P_s \mu'_s$, and by path independence, $z \in Ch_{d_1}(\mu'_{d_1} \cup$ $\{z\}$). Furthermore, since student *s* is matched with district d_1 in μ' , $d(s) = d_1$. Therefore, $\widetilde{Ch}_{d_1}(\mu'_{d_1} \cup \{z\}) = \widetilde{Ch}'_{d_1}(\mu'_{d_1} \cup \{z\})$ by definition of \widetilde{Ch}'_{d_1} and construction of μ' . Hence, $z \in \widetilde{Ch}'_{d_1}(\mu'_{d_1} \cup \{z\})$, which contradicts the fact that μ' is stable under $(\widetilde{Ch}'_d)_{d \in \mathcal{D}}$. Hence, $\mu'_{d_1} \cup \mu^1_{d_1}$ is feasible for students. Feasibility for students implies that $Ch_{d_1}(\mu'_{d_1} \cup \mu^1_{d_1}) \supseteq$ $\widetilde{Ch}'_{d_1}(\mu'_{d_1} \cup \mu^1_{d_1})$. Path independence and construction of $\mu^1_{d_1}$ yield $\mu^1_{d_1} = \widetilde{Ch}_{d_1}(\mu'_{d_1} \cup \mu^1_{d_1})$. Furthermore, there exists no student *s*, such that $d(s) = d_1$, who has a contract in $\mu_{d_1}^1 \setminus \mu'$, as this would contradict stability of μ' under $(\widetilde{Ch}'_d)_{d\in\mathcal{D}}$. This implies, by definition of \widetilde{Ch}'_{d_1} , that $\widetilde{Ch}'_{d_1}(\mu'_{d_1} \cup \mu^1_{d_1}) = \widetilde{Ch}'_{d_1}(\mu'_{d_1})$, and, by stability of μ' under $(\widetilde{Ch}'_{d_1})_{d \in \mathcal{D}}, \widetilde{Ch}'_{d_1}(\mu'_{d_1}) = \mu'_{d_1}$. Therefore, $\mu_{d_1}^1 = \widetilde{Ch}_{d_1}(\mu'_{d_1} \cup \mu_{d_1}^1) \supseteq \widetilde{Ch}'_{d_1}(\mu'_{d_1} \cup \mu_{d_1}^1) = \mu'_{d_1} = \mu_{d_1}^0$, which means that district d_1 does not reject any contracts.

Now consider district d_n where n > 1. There are two cases to consider. First consider the case when $d_n \neq d_i$ for every i < n. In this case, $\mu_{d_n}^{n-1} \subseteq \mu_{d_n}^0 = \mu'_{d_n}$. We repeat the same arguments as in the previous paragraph. Stability of μ' under $(\widetilde{Ch}'_d)_{d\in\mathcal{D}}$ and path independence of \widetilde{Ch}'_{d_n} implies that $\mu_{d_n}^n \cup \mu_{d_n}^{n-1}$ is feasible for students. Therefore, $\widetilde{Ch}_{d_n}(\mu_{d_n}^{n-1} \cup \mu_{d_n}^n) \supseteq \widetilde{Ch}'_{d_n}(\mu_{d_n}^{n-1} \cup \mu_{d_n}^n)$. Furthermore, there exists no student *s*, such that $d(s) = d_n$, who has a contract in $\mu_{d_n}^n \setminus \mu_{d_n}^{n-1}$. As a result, by definition of \widetilde{Ch}'_{d_n} and by path independence, $\widetilde{Ch}'_{d_n}(\mu_{d_n}^{n-1} \cup \mu_{d_n}^n) = \widetilde{Ch}'_{d_n}(\mu_{d_n}^{n-1}) = \mu_{d_n}^{n-1}$. As in the previous paragraph, we conclude that $\mu_{d_n}^n = \widetilde{Ch}_{d_n}(\mu_{d_n}^{n-1} \cup \mu_{d_n}^n) \supseteq \widetilde{Ch}'_{d_n}(\mu_{d_n}^{n-1} \cup \mu_{d_n}^n) = \mu_{d_n}^{n-1}$.

The second case is when there exists i < n such that $d_i = d_n$. Let i^* be the last such step before n. Since student welfare improves at every step before n by the mathematical induction hypothesis, $\mu_{d_n}^{i^*-1} \cup X_{d_n}^{i^*} \supseteq \mu_{d_n}^{n-1} \cup X_{d_n}^n$. By definition, $\mu_{d_n}^{i^*} = \widetilde{Ch}_{d_n}(\mu_{d_n}^{i^*-1} \cup X_{d_n}^{i^*})$, which implies by path independence that $\mu_{d_n}^{n-1} \subseteq \widetilde{Ch}_{d_n}(\mu_{d_n}^{n-1} \cup X_{d_n}^n) = \mu_{d_n}^n$ since $\mu_{d_n}^{n-1} \subseteq \mu_{d_n}^{i^*}$.

Finally, we need to show that the improvement algorithm terminates. We claim that $\mu_{d_n}^n \neq \mu_{d_n}^{n-1}$. Suppose, for contradiction, that these two matchings are the same. Then, by path independence of \widetilde{Ch}_{d_n} , for every $x \in X_{d_n}^n$, $\widetilde{Ch}_{d_n}(\mu_{d_n}^{n-1} \cup \{x\}) = \mu_{d_n}^{n-1}$. This is a contradiction because there exists at least one blocking contract associated with district d_n .

Therefore, district d_n gets at least one new contract at Step n. Hence, at least one student gets a strictly more preferred contract at every step of the algorithm while every other student gets a weakly more preferred contract. Since the number of contracts is finite, the algorithm has to end in a finite number of steps.

Because the interdistrict SPDA outcome under $(Ch_d)_{d\in\mathcal{D}}$ is the same as the interdistrict SPDA outcome under $(\widetilde{Ch}_d)_{d\in\mathcal{D}}$ and the interdistrict SPDA outcome under $(Ch'_d)_{d\in\mathcal{D}}$ is the same as the interdistrict SPDA outcome under $(\widetilde{Ch}')_{d\in\mathcal{D}}$, the lemma implies that every student weakly prefers the outcome of interdistrict SPDA under $(Ch_d)_{d\in\mathcal{D}}$ to the outcome of intradistrict SPDA (which is the same as the interdistrict SPDA outcome under $(Ch'_d)_{d\in\mathcal{D}}$). This completes the proof of the first part.

To prove the second part of the theorem, we show that if at least one district's admissions rule fails to favor own students, then there exists a preference profile such that not every student is weakly better off under interdistrict SPDA. Suppose that for some district d, there exists a matching X, which is feasible for students, such that $Ch_d(X)$ is not a superset of $Ch_d(X^*)$, where $X^* \equiv \{x \in X | d(s(x)) = d\}$. Now, consider a matching Y where (i) all students from district d are matched with schools in district d, (ii) Y is feasible, and (iii) $Y \supseteq Ch_d(X^*)$. The existence of such a Y follows from the fact that $Ch_d(X^*)$ is feasible and $k_{d'} \leq \sum_{c:d(c)=d'} q_c$, for every district d' (that is, there are enough seats in district d' to match all students from district d'.) Because Y is feasible and Ch_d is acceptant, $Ch_d(Y_d) = Y_d$.

Now consider the following student preferences. First we consider students from district *d*. Each student *s* who has a contract in X^* ranks X_s^* as her top choice. Note that doing so is well defined because X^* is feasible for students. Each student *s* who has a contract in $X^* \setminus Ch_d(X^*)$ ranks contract Y_s as her second top choice. Note that, in this case, Y_s cannot be the same as X_s^* because $Ch_d(Y_d) = Y_d$ and Ch_d is path independent. Each student *s* who has a contract in $Y \setminus X^*$ ranks that contract as her top choice. Next we consider students from the other districts. Each student *s* who has a contract in $X \setminus X^*$ ranks that contract as her top choice. Any other student ranks a contract not associated with district *d* as her top choice. Complete the rest of the student preferences arbitrarily.

Consider SPDA for district d for intradistrict school choice. At the first step, students who have a contract in X^* propose that contract. The remaining students that have contracts in $Y \setminus X^*$ propose the associated contracts. Because Y is feasible, Y contains $Ch_d(X^*)$, and Ch_d is acceptant, only contracts in $X^* \setminus Ch_d(X^*)$ are rejected. At the second step, these students propose their contracts in Y_d , and the set of proposals that the district considers is Y_d . Because $Ch_d(Y_d) = Y_d$, no contract is rejected, and SPDA stops and returns Y_d . In particular, every student who has a contract in $Ch_d(X^*)$ has the corresponding contract at the outcome.

In interdistrict SPDA, at the first step, each student who has a contract in X proposes that contract and every other student proposes a contract associated with a district different from d. District d considers X (or X_d), and tentatively accepts $Ch_d(X)$. Because $Ch_d(X) \not\supseteq Ch_d(X^*)$ by assumption, at least one student who has a contract in $Ch_d(X^*)$ is rejected. Therefore, this student is strictly worse off under interdistrict school choice.

Proof of Theorem 9. To show the result, we first introduce the following weakening of the substitutability condition (Hatfield and Kojima, 2008). A district admissions rule Ch_d satisfies *weak substitutability* if, for every $x \in X \subseteq Y \subseteq \mathcal{X}$ with $x \in Ch_d(Y)$ and $|Y_s| \leq 1$ for each $s \in S$, it must be that $x \in Ch_d(X)$.

Under weak substitutability, the following result is known (the statement is slightly modified for the present setting).

Theorem 10 (Hatfield and Kojima (2008)). Let d and d' be two distinct districts. Suppose that Ch_d satisfies IRC but violates weak substitutability. Then, there exist student preferences and a path-independent admissions rule for d' such that, regardless of the other districts' admissions rules, no stable matching exists.

Given this result, for our purposes it suffices to show the following.

Theorem 9'. Let *d* be a district. There exist a set of students, their types, schools in *d*, and typespecific ceilings for *d* such that there is no district admissions rule of *d* that has district-level typespecific ceilings, is *d*-weakly acceptant, and satisfies IRC and weak substitutability.

To show this result, consider a district d with $k_d = 2$. There are three schools c_1 , c_2 , c_3 in the district, each with capacity one, and four students s_1 , s_2 , s_3 , s_4 of which two are from a different district. Students s_1 and s_2 are of type t_1 and students s_3 and s_4 are of type t_2 . The district-level type-specific ceilings are as follows: $q_d^{t_1} = q_d^{t_2} = 1$.

Suppose, for contradiction, that the district admissions rule has district-level typespecific ceilings, is *d*-weakly acceptant, and satisfies IRC and weak substitutability.

Consider $Ch_d(\{(s_1, c_1), (s_2, c_1), (s_3, c_1), (s_4, c_1)\})$. Since types are symmetric and two students are symmetric within each type, without loss of generality, we can assume $Ch_d(\{(s_1, c_1), (s_2, c_1), (s_3, c_1), (s_4, c_1)\}) = \{(s_1, c_1)\}.$

Next, consider $Ch_d(\{(s_2, c_2), (s_3, c_2), (s_4, c_2)\})$. This can be either equal to $\{(s_2, c_2)\}$ or $\{(s_3, c_2)\}$ (the case when it is equal to $\{(s_4, c_2)\}$ is symmetric to the case when $\{(s_3, c_2)\}$. We analyze these two cases separately.

(1) Suppose $Ch_d(\{(s_2, c_2), (s_3, c_2), (s_4, c_2)\}) = \{(s_2, c_2)\}$. Then, by IRC, we conclude that $Ch_d(\{(s_2, c_2), (s_3, c_2)\}) = \{(s_2, c_2)\}$. Next, we argue that $Ch_d(\{(s_1, c_1), (s_2, c_2), (s_3, c_2)\}) = \{(s_2, c_2)\}$. This is because the only two

cases that satisfy d-weak acceptance and type-specific ceilings are $\{(s_2, c_2)\}$ and $\{(s_1, c_1), (s_3, c_2)\}$. The latter would violate weak substitutability since in that case (s_3, c_2) would be accepted in a larger set $\{(s_1, c_1), (s_2, c_2), (s_3, c_2)\}$ and rejected from a smaller set $\{(s_2, c_2), (s_3, c_2)\}$. Then, by IRC, $Ch_d(\{(s_1, c_1), (s_2, c_2), (s_3, c_2)\}) = \{(s_2, c_2)\}$ implies $Ch_d(\{(s_1, c_1), (s_2, c_2), (s_3, c_1)\}) = \{(s_2, c_2), (s_3, c_1)\}$ more that $Ch_d(\{(s_1, c_1), (s_2, c_2), (s_3, c_1)\}) = \{(s_2, c_2), (s_3, c_1)\}$ since by weak substitutability (s_1, c_1) cannot be chosen, and therefore (s_2, c_2) and (s_3, c_1) have to be chosen due to *d*-weak acceptance. Next, again by weak substitutability, we note that $Ch_d(\{(s_1, c_1), (s_2, c_2), (s_3, c_1)\}) = \{(s_3, c_1)\}$. Finally, we note that this contradicts with $Ch_d(\{(s_1, c_1), (s_2, c_1), (s_3, c_1), (s_4, c_1)\}) = \{(s_1, c_1)\}$ and IRC.

- (2) Suppose $Ch_d(\{(s_2, c_2), (s_3, c_2), (s_4, c_2)\}) = \{(s_3, c_2)\}$. Consider $Ch_d(\{(s_2, c_3), (s_4, c_3)\})$. This can be either $\{(s_2, c_3)\}$ or $\{(s_4, c_3)\}$. We consider these two possible cases separately. These two subcases will follow similar arguments to Case (1) above and change the indexes appropriately in order to get a contradiction.
 - (a) Suppose $Ch_d(\{(s_2, c_3), (s_4, c_3)\}) = \{(s_2, c_3)\}$. Next, we argue that $Ch_d(\{(s_1, c_1), (s_2, c_3), (s_4, c_3)\}) = \{(s_2, c_3)\}$. This is because the only two cases that satisfy d-weak acceptance and type-specific ceilings are $\{(s_2, c_3)\}$ and $\{(s_1, c_1), (s_4, c_3)\}$. The latter would violate weak substitutability since in that case (s_4, c_3) would be accepted in a larger set $\{(s_1, c_1), (s_2, c_3), (s_4, c_3)\}$ and rejected from a smaller set $\{(s_2, c_3), (s_4, c_3)\}$. Then, by IRC, $Ch_d(\{(s_1, c_1), (s_2, c_3), (s_4, c_3)\}) = \{(s_2, c_3)\}$ implies $Ch_d(\{(s_1, c_1), (s_2, c_3), (s_4, c_3)\}) = \{(s_2, c_3), (s_4, c_1)\}\}$ since by weak substitutability (s_1, c_1) cannot to be chosen, therefore (s_2, c_3) and (s_4, c_1) have to be chosen due to d-weak acceptance. Next, again by weak substitutability, we note that $Ch_d(\{(s_1, c_1), (s_2, c_3), (s_4, c_1)\}) = \{(s_2, c_3), (s_4, c_1)\}$ implies $Ch_d(\{(s_1, c_1), (s_4, c_1)\}) = \{(s_4, c_1)\}$. Finally, we note that this contradicts with $Ch_d(\{(s_1, c_1), (s_2, c_1), (s_3, c_1), (s_4, c_1)\}) = \{(s_1, c_1)\}$ and IRC.
 - (b) Suppose $Ch_d(\{(s_2, c_3), (s_4, c_3)\}) = \{(s_4, c_3)\}$. Next, we argue that $Ch_d(\{(s_2, c_3), (s_3, c_2), (s_4, c_3)\}) = \{(s_4, c_3)\}$. This is because the only two cases that satisfy d-weak acceptance and type-specific ceilings are $\{(s_4, c_3)\}$ and $\{(s_2, c_3), (s_3, c_2)\}$. The latter would violate weak substitutability since in that case (s_2, c_3) would be accepted in a larger set $\{(s_2, c_3), (s_3, c_2), (s_4, c_3)\}$ and rejected from a smaller set $\{(s_2, c_3), (s_4, c_3)\}$. Then, by IRC, $Ch_d(\{(s_2, c_3), (s_3, c_2), (s_4, c_3)\}) = \{(s_4, c_3)\}$ implies $Ch_d(\{(s_3, c_2), (s_4, c_3)\}) = \{(s_4, c_3)\}$. Then we note that $Ch_d(\{(s_2, c_2), (s_3, c_2), (s_4, c_3)\}) = \{(s_2, c_2), (s_4, c_3)\}$

since by weak substitutability (s_3, c_2) cannot to be chosen, therefore (s_4, c_3) and (s_2, c_2) have to be chosen due to d-weak acceptance. Next, again by weak substitutability, we note that $Ch_d(\{(s_2, c_2), (s_3, c_2), (s_4, c_3)\}) = \{(s_2, c_2), (s_4, c_3)\}$ implies $Ch_d(\{(s_2, c_2), (s_3, c_2)\}) = \{(s_2, c_2)\}$. Finally, we note that this contradicts with $Ch_d(\{(s_2, c_2), (s_3, c_2), (s_4, c_2)\}) = \{(s_3, c_2)\}$ and IRC.